

CONTINUOUS FAMILIES OF HAMILTONIAN TORUS ACTIONS

ANDRÉS VIÑA

ABSTRACT. We determine conditions under which two Hamiltonian torus actions on a symplectic manifold M are homotopic by a family of Hamiltonian torus actions, when M is a toric manifold and when M is a coadjoint orbit.

MSC 2000: 53D05, 57S05

1. INTRODUCTION

Let (M, ω) be a closed symplectic $2n$ -manifold, and let G be a compact Lie group. Given two Hamiltonian G -actions on M , it is a natural issue to analyze if one can be deformed in the another by a continuous family of Hamiltonian G -actions. In this paper we will consider this matter; that is, we will determine necessary conditions for two Hamiltonian G -actions on M to be connected by a homotopy consisting of Hamiltonian G -actions.

The mathematical setting is the following: $\text{Ham}(M)$ will be the Hamiltonian group of (M, ω) [20], [21], and $\text{Hom}(G, \text{Ham}(M))$ will denote the space of all Lie group homomorphisms ψ from G to $\text{Ham}(M)$; that is, the space consisting of the Hamiltonian G -actions on M .

Given $\psi, \psi' \in \text{Hom}(G, \text{Ham}(M))$, we write $\psi \sim \psi'$ iff ψ and ψ' belong to the same connected component of $\text{Hom}(G, \text{Ham}(M))$. In other words, $\psi \sim \psi'$ iff there is a continuous family $\{\psi^s : G \rightarrow \text{Ham}(M)\}_{s \in [0,1]}$ of Lie group homomorphisms, such that $\psi^0 = \psi$ and $\psi^1 = \psi'$; that is, ψ and ψ' are homotopic by a family of *group homomorphisms*. The corresponding quotient space is denoted by $[G, \text{Ham}(M)]_{gh}$.

In this paper we will consider the following three general aspects:

1. *Necessary conditions for the $\sim$ -equivalence.* It would be desirable to obtain necessary and sufficient conditions for the $\sim$ -equivalence of two G -actions in the general case, but we will prove more modest results. To each Hamiltonian G -action ψ we will associate a cohomology class S_ψ , such that the map S_- is compatible with the relation $\sim$, and we will use the map S_- to distinguish different elements of $[G, \text{Ham}(M)]_{gh}$ (see Theorem 1). In some particular cases we will give direct proofs of the $\sim$ -equivalence of two actions on which the map S_- takes the same value.

2. *Equivalence relations between toric actions.* A G -action can be regarded as a representation of G in the group $\text{Ham}(M)$, so we can consider another natural equivalence relation in the space $\text{Hom}(G, \text{Ham}(M))$: Two Hamiltonian G -actions on M , ψ and ψ' , are said to be *r-equivalent* if there exist an element $h \in \text{Ham}(M)$, such that $h \circ \psi_g \circ h^{-1} = \psi'_g$ for all $g \in G$; that is, if they are equivalent as

Key words and phrases. Hamiltonian diffeomorphisms, toric actions, coadjoint orbits.

This work has been partially supported by Ministerio de Ciencia y Tecnología, grant MAT2003-09243-C02-01.

representations of G in the group $\text{Ham}(M)$. We will compare the r -equivalence and the relation $\sim$ in the space of toric actions on M . We will prove, among other properties, that the r -equivalence implies the $\sim$ -equivalence, and that the equality of the moment polytopes of ψ and ψ' is a necessary condition for the $\sim$ -equivalence of them (see Theorem 3).

3. *$\sim$ -equivalence of $U(1)$ -actions.* When $G = U(1)$, each $\psi \in \text{Hom}(G, \text{Ham}(M))$ determines an element of $\{\psi\} \in \pi_1(\text{Ham}(M))$. We will compare our relation $\sim$ with the equivalence in $\pi_1(\text{Ham}(M))$ in some particular cases for which the homotopy type of $\text{Ham}(M)$ is known. In [8] Gromov proved that $\text{Ham}(\mathbb{C}P^2, \omega_{FS})$ is homotopy equivalent to the group $PU(3)$, where ω_{FS} is the Fubini-Study symplectic structure, and in [1] Abreu and McDuff determined the homology of the group of symplectomorphisms of a rational ruled surface. In order to carry out the aforementioned comparison and also by its very interest, we will study the relation $\sim$ between $U(1)$ -actions in two cases:

a) When M is the total space of the fibration

$$(1.1) \quad \mathbb{P}(L_1 \oplus \cdots \oplus L_{n-1} \oplus \mathbb{C}) \rightarrow \mathbb{C}P^1,$$

L_i being a holomorphic line bundle over $\mathbb{C}P^1$.

b) When the manifold M is a coadjoint orbit; we look more closely at the case of $U(1)$ -actions on coadjoint orbits of $SU(n)$ diffeomorphic to Grassmann manifolds.

The comparison of the results of Gromov and Abreu-McDuff above mentioned and our results shows the non-injectivity of the obvious map

$$(1.2) \quad [\psi] \in [U(1), \text{Ham}(M)]_{gh} \mapsto \{\psi\} \in \pi_1(\text{Ham}(M)),$$

when M is a Hirzebruch surface and when $M = \mathbb{C}P^2$ (see Remark below Theorem 7, the paragraph after Theorem 8 and Remark below Theorem 11).

Statement of main results

1. Necessary conditions for the $\sim$ -equivalence. An action ψ of G on M determines the fibration $p : M_\psi := EG \times_G M \rightarrow BG$, with fibre M , on the classifying space BG of the group G . If ψ is Hamiltonian, the fibration M_ψ supports the coupling class $c_\psi \in H^2(M_\psi, \mathbb{R})$ [20], [10]. Then fibre integration of $\exp(c_\psi)$ defines an element S_ψ in the cohomology $H(BG)$. We will prove that $S_\psi = S_{\psi'}$, if the G -actions ψ and ψ' are $\sim$ -equivalent. That is,

Theorem 1. *Let ψ and ψ' be two Hamiltonian G -actions on M , if $S_\psi \neq S_{\psi'}$, then $[\psi] \neq [\psi'] \in [G, \text{Ham}(M)]_{gh}$.*

The cohomology class S_ψ can be calculated by the localization formula of the G -equivariant cohomology. This formula gives an expression for S_ψ which is a linear combination of exponentials functions. If G is the m -dimensional torus, the coefficients of this combination are rational functions of m variables $(u_1, \dots, u_m)$. On the other hand, given real constants γ_{ij} , the exponential functions $F_i(u_1, \dots, u_m) := \exp(\sum_j \gamma_{ij} u_j)$ are $\mathbb{R}(u_1, \dots, u_m)$ -linearly independent (see Lemma 14). This property will allow us to compare the values of the map S_- on different G -actions. The comparison of the values taken by this map, together with Theorem 1, will permit us to determine necessary conditions, with a simple geometric content, for two torus actions to be $\sim$ -equivalent (see, for example, Theorem 2 and Theorem 5).

Let us assume that (M, ω) is equipped with a Hamiltonian action of the Lie group G . If $X \in \mathfrak{g}$ defines a circle action ψ^X on M , we say that X is of type

s when a connected component of the fixed point set of ψ^X is a $2s$ -dimensional submanifold of M , and there is no submanifold of greater dimension in the fixed point set. The set of elements of type s will be denoted $\text{Type}(s)$. We will prove the following result, which gives a necessary condition for the $\sim$ -equivalence whose geometric meaning is clear

Theorem 2. *If $X, Y \in \mathfrak{g}$ are of different type, then the corresponding circle actions ψ^X and ψ^Y are not homotopic by means of a family of $U(1)$ -actions.*

2. Equivalence relations between torus actions.


In Section 2 we will consider *toric actions* on M ; that is, Hamiltonian effective actions of the n -dimensional torus $T := U(1)^n$ on M . A toric action ψ defines a moment polytope $\Delta(\psi) = \Delta$ in $\mathfrak{t}^*$, and this polytope is uniquely determined by ψ up to translation.

Let ψ, ψ' be toric actions on M , and let Δ, Δ' denote the moment polytopes with center of mass at 0 associated to ψ and ψ' , respectively. By μ and μ' we denote the corresponding moment maps, such that $\text{im } \mu = \Delta$ and $\text{im } \mu' = \Delta'$. We state the following properties:

- (i) $\Delta = \Delta'$.
- (ii) $\Delta = \Delta'$ and there exists $h \in \text{Symp}_0(M)$ such that $h \circ \psi_g \circ h^{-1} = \psi'_g$ for all $g \in T$, where $\text{Symp}_0(M)$ is the connected component of the identity in the group $\text{Symp}(M)$ of symplectomorphisms of M .
- (iii) ψ and ψ' are r -equivalent.
- (iv) $S_\psi = S_{\psi'}$.
- (v) $[\psi] = [\psi'] \in [T, \text{Ham}(M)]_{gh}$.

We will prove the following Theorem

Theorem 3. *The above properties satisfy the implications described in the following diagram*



From this theorem it follows, among other properties, that the maps $S_\cdot$ and $\Delta(\cdot)$ are equivalent tools for distinguishing $\sim$ -inequivalent toric actions. The equality $\Delta = \Delta'$ is a very simple geometric necessary condition for the equality $[\psi] = [\psi']$, and a sufficient condition can be obtained by complementing that necessary condition as it is stated in (ii), but this complementary condition is not easy to check, in general.


Given $\varphi, \varphi' \in \text{Hom}(G, \text{Ham}(M))$, we write $\varphi \sim_{rp} \varphi'$ if φ and φ' are $\sim$ -equivalent “up to reparametrization” of G ; that is, if there is an automorphism v of G such that $\varphi \circ v \sim \varphi'$. Following Karshon [15] we say that the Hamiltonian actions φ and φ' are *conjugate* if they are representations in $\text{Symp}(M)$ equivalent “up to reparametrization”; that is, if there is $v \in \text{Aut}(G)$ and $\xi \in \text{Symp}(M)$ such that $\xi \circ (\varphi \circ v)(g) \circ \xi^{-1} = \varphi'(g)$, for all $g \in G$. To complete the scheme we define the r -equivalence up to reparametrization: φ and φ' are r -equivalent up to reparametrization iff there exist $v \in \text{Aut}(G)$ and $h \in \text{Ham}(M)$ such that $h \circ (\varphi(v(g))) \circ h^{-1} = \varphi'(g)$, for all $g \in G$.

Some of the implications in Theorem 3 remains valid when the equivalence relations are replaced by the corresponding “up to reparametrization” relations. For ψ and ψ' toric actions on M we state the following properties:

- (a) ψ and ψ' are r -equivalent up to reparametrization.
- (b) $\psi \sim_{rp} \psi'$.
- (c) ψ and ψ' are conjugate.
- (d) There exists $V \in GL(n, \mathbb{Z})$, such that $V(\Delta) = \Delta'$.

The following Proposition is also proved in Section 2

Proposition 4. *For the above properties the implications of the following diagram*



hold.

Let M be a toric manifold and ψ the effective action of T which endows M with the toric structure. If X belongs to the integer lattice of $\mathfrak{t}$, it generates an S^1 -action ψ^X . If the element X is of type s , then there is an s -face f in the moment polytope Δ such that $\langle f, X \rangle = 0$, but there is no an r -face orthogonal to X with $r > s$. If X is of Type(0), the fixed point set for ψ^X is the set $\{\mu^{-1}(P_i) \mid i = 1, \dots, s\}$, where the P_i 's are the vertices of the moment polytope Δ and μ the moment map.

We will prove the following Theorem that gives a simple geometric condition, under which two elements X, Y of type 0 define circle actions ψ^X and ψ^Y no homotopic by a family of $U(1)$ -actions.

Theorem 5. *Let M be a toric manifold and Δ the moment polytope with center of mass at 0. If X, Y belong to the integer lattice of $\mathfrak{t}$, they are in Type(0) and*

$$\{\langle P_i, X \rangle : i = 1, \dots, s\} \neq \{\langle P_i, Y \rangle : i = 1, \dots, s\},$$

$P_1, \dots, P_s$ being the vertices of Δ . Then $[\psi^X] \neq [\psi^Y] \in [U(1), \text{Ham}(M)]_{gh}$.

Proposition 21 in Section 2 is the version “up to reparametrization” of Theorem 5.

3. $\sim$ -equivalence of $U(1)$ -actions.

S^1 -actions on $\mathbb{P}(L_1 \oplus \dots \oplus L_{n-1} \oplus \mathbb{C})$.

If Δ is a Delzant polytope in $(\mathbb{R}^n)^*$, we denote by M_Δ the symplectic toric manifold associated to Δ [5], [9]. We will study the relation $\sim$ when M_Δ is diffeomorphic to the total space of the fibration (1.1). In this case the result stated in Theorem 5 will be completed giving a sufficient condition for two elements in the integer lattice to define $\sim$ -equivalent $U(1)$ -actions.

Let us assume that M_Δ is diffeomorphic to $\mathbb{P}(L_1 \oplus \dots \oplus L_{n-1} \oplus \mathbb{C})$, where L_i is the holomorphic line bundle over $\mathbb{C}P^1$ with first Chern number $-a_i$. The S^1 -action defined by $\mathbf{b} \in \mathbb{Z}^n$ is denoted by $\psi_{\mathbf{b}}$. To study the relation $\sim$ between the $\psi_{\mathbf{b}}$'s we introduce the following definition:

Given $\mathbf{b} = (b_1, \dots, b_n)$ and $\mathbf{b}' = (b'_1, \dots, b'_n)$ two elements of the lattice $\mathbb{Z}^n$. We write $\mathbf{b} \equiv \mathbf{b}'$ iff either $\mathbf{b} = \mathbf{b}'$, or

$$b'_j = b_j + a_j b_n, \text{ for } j = 1, \dots, n-1 \text{ and } b'_n = -b_n.$$

We will prove the following Theorem, which gives the aforesaid sufficient condition for the $\sim$ -equivalence between $\psi_{\mathbf{b}}$ and $\psi_{\mathbf{b}'}$.

Theorem 6. *Let $\mathbf{b}$ and $\mathbf{b}'$ be two elements of the integer lattice $\mathbb{Z}^n$. Let $\psi_{\mathbf{b}}$, $\psi_{\mathbf{b}'}$ be the respective circle actions on the symplectic manifold M_{Δ} diffeomorphic to $\mathbb{P}(L_1 \oplus \cdots \oplus L_{n-1} \oplus \mathbb{C})$. If $\mathbf{b} \equiv \mathbf{b}'$, then $\psi_{\mathbf{b}} \sim \psi_{\mathbf{b}'}$.*

When the manifold M_{Δ} is a Hirzebruch surface we will prove a stronger result than the one stated in Theorem 6. Let M_{Δ} be the Hirzebruch surface $\mathbb{P}(L_k \oplus \mathbb{C})$ (where L_k is the holomorphic line bundle over $\mathbb{C}P^1$ with first Chern number k). If $\mathbf{b} = (b_1, b_2)$ is an element in $\mathbb{Z}^2$, set $S_{\mathbf{b}} = S_{\psi_{\mathbf{b}}}$. We will prove the following Theorem

Theorem 7. *Let M_{Δ} be the Hirzebruch surface $\mathbb{P}(L_k \oplus \mathbb{C})$, with $k \neq 0$, and $\mathbf{b}, \mathbf{b}' \in \mathbb{Z}^2$, then the following statements are equivalent*

- (a) $\mathbf{b} \equiv \mathbf{b}'$.
- (b) $\psi_{\mathbf{b}} \sim \psi_{\mathbf{b}'}$.
- (c) $S_{\mathbf{b}} = S_{\mathbf{b}'}$.

Remark. Set $\mathbf{c} = (1, 0)$ and $\mathbf{c}' = (0, 1)$. When k is odd,

$$\{\psi_{\mathbf{c}}\}, \{\psi_{\mathbf{c}'}\} \in \pi_1(\text{Ham}(\mathbb{P}(L_k \oplus \mathbb{C})))$$

are elements of infinite order in this group (see Section 4 of [22]). On the other hand, $\pi_1(\text{Ham}(\mathbb{P}(L_k \oplus \mathbb{C}))) = \mathbb{Z}$ (see [1]). Hence there exist $m, n \in \mathbb{Z} \setminus \{0\}$ such that $\{\psi_{m\mathbf{c}}\} = \{\psi_{n\mathbf{c}'}\}$. As the condition (a) of Theorem 7 does not hold for $m\mathbf{c}$ and $n\mathbf{c}'$, it follows that $\psi_{m\mathbf{c}}$ and $\psi_{n\mathbf{c}'}$ are not $\sim$ -equivalent; that is, the map (1.2) is not injective for the Hirzebruch surface we are considering.

When $k = 0$ the manifold $\mathbb{P}(L_k \oplus \mathbb{C})$ is diffeomorphic to $S^2 \times S^2$. In this case let τ and σ denote the total areas of respective the spheres in the product $S^2 \times S^2$. We will prove following result

Theorem 8. *If $b_1 b_2 \neq 0 \neq b'_1 b'_2$ and $\sigma/\tau \notin \mathbb{Q}$, then the following statements are equivalent*

- (i) $|b'_1| = |b_1|$ and $|b'_2| = |b_2|$.
- (ii) $S_{\mathbf{b}} = S_{\mathbf{b}'}$.
- (iii) $\psi_{\mathbf{b}} \sim \psi_{\mathbf{b}'}$.

When $\sigma > 1$ and $\tau = 1$,

$$(1.3) \quad \pi_1(\text{Ham}(S^2 \times S^2)) = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}.$$

The first two summands of the right hand of (1.3) come from the action of $SO(3) \times SO(3)$ on $S^2 \times S^2$. The generator of the third summand may be represented by the loop $\{R^t\}_t$, where

$$R^t : (u, v) \in S^2 \times S^2 \mapsto (u, R_u^t(v)) \in S^2 \times S^2,$$

R_u^t being the rotation of the fiber S^2 , with angle $2\pi t$ and axis through u (see [1], [19]). Therefore $\psi_{\mathbf{c}}$, with $\mathbf{c} = (1, 0)$, is a representative of the generator of the first summand in the right hand of (1.3). Thus $\{\psi_{\mathbf{c}}\} = \{\psi_{m\mathbf{c}}\} \in \pi_1(\text{Ham}(S^2 \times S^2))$, for all $m \in 2\mathbb{Z} + 1$. But, according to Theorem 8, from

$$[\psi_{\mathbf{c}}] = [\psi_{m\mathbf{c}}] \in [U(1), \text{Ham}(S^2 \times S^2)]_{gh}$$

it follows $m = \pm 1$, when $\sigma \notin \mathbb{Q}$.

Although the results stated in Theorem 7 and in Theorem 8, the equality $S_{\mathbf{b}} = S_{\mathbf{b}'}$ is not equivalent to $[\psi_{\mathbf{b}}] = [\psi_{\mathbf{b}'}] \in [U(1), \text{Ham}(M)]_{gh}$, in general. As we said, $S_{\mathbf{b}} = S_{\mathbf{b}'}$ is a necessary condition for the equality of the classes $[\psi_{\mathbf{b}}]$ and $[\psi_{\mathbf{b}'}]$. But if (M, ω) possesses certain symmetries it can happen that $S_{\mathbf{b}} = S_{\mathbf{b}'}$ and $[\psi_{\mathbf{b}}] \neq [\psi_{\mathbf{b}'}]$ (see Remark in Subsection 3.2).

Circle actions on coadjoint orbits.

If G is a compact connected Lie group and $\eta \in \mathfrak{g}^*$, the stabilizer of η for the coadjoint action contains a maximal torus H . The integer elements of $\mathfrak{h}$ define Hamiltonian circle actions on the coadjoint orbit $\mathcal{O}$ of η . We will consider the equivalence relation $\sim$ in the set of those S^1 -actions on the orbit $\mathcal{O}$. By W we denote the Weyl group of the pair (G, H) . We will prove the following Theorem

Theorem 9. *Let X, X' be two elements of the integer lattice of $\mathfrak{h}$. If there exists $w \in W$ such that $w(X) = X'$, then the circle actions defined on $\mathcal{O}$ by X and X' are $\sim$ -equivalent.*

A particular case is when $G = SU(n)$, H its diagonal subgroup and $\mathcal{O}$ is the orbit of an element whose stabilizer contains H . In this case we have the following Corollary

Corollary 10. *If $X = (x_1, \dots, x_n)$, $X' = (x'_1, \dots, x'_n) \in (2\pi i\mathbb{Z})^n$ and there exists a permutation τ of the set $\{1, \dots, n\}$, such that $x'_j = x_{\tau(j)}$ for all j , then the circle actions defined by X and X' on the orbit $\mathcal{O}$ of $SU(n)$ are $\sim$ -equivalent.*

When the coadjoint orbit is diffeomorphic to $\mathbb{C}P^{n-1}$ we will prove a stronger result.

Theorem 11. *Let $\mathcal{O}$ be a coadjoint orbit of $SU(n)$ diffeomorphic to $\mathbb{C}P^{n-1}$, and X, X' diagonal elements of $\mathfrak{su}(n)$ such that $X, X' \in (2\pi i\mathbb{Z})^n$. If ψ and ψ' denote the circle actions on $\mathcal{O}$ defined by X and X' respectively. Then the following are equivalent*

- (a) $[\psi] = [\psi'] \in [U(1), \text{Ham}(\mathcal{O})]_{gh}$.
- (b) $S_{\psi} = S_{\psi'}$.
- (c) *There exists a permutation τ of the set $\{1, \dots, n\}$, such that $x'_j = x_{\tau(j)}$ for $j = 1, \dots, n$.*

Remark. By the preceding Theorem, the elements $X \in (2\pi i\mathbb{Z})^3$ define an infinite family in $[U(1), \text{Ham}(\mathbb{C}P^2)]_{gh}$. But in contrast $\sharp(\pi_1(\text{Ham}(\mathbb{C}P^2))) = 3$, since $\text{Ham}(\mathbb{C}P^2)$ has the homotopy type of $PU(3)$ [8]. Thus the map (1.2) is not injective when $M = \mathbb{C}P^2$.

To state a necessary condition for the equality $[\psi] = [\psi']$ when the coadjoint orbit of $SU(n)$ is the Grassmann manifold $G_k(\mathbb{C}^n)$ (of the k -subspaces in $\mathbb{C}^n$) we need to introduce new notations. By $\mathcal{C}_k$ we denote the set of combinations of the set $\{1, \dots, n\}$ taken k at a time. If $X = (ia_1, \dots, ia_n)$ and $\mathfrak{d} = \{j_1, \dots, j_k\} \in \mathcal{C}_k$ we write $\alpha_{\mathfrak{d}} = \sum_{l=1}^k a_{j_l}$. We will prove the following result

Theorem 12. *Let $\mathcal{O}$ be a coadjoint orbit of $SU(n)$ diffeomorphic to the Grassmann manifold $G_k(\mathbb{C}^n)$, and X, X' diagonal regular elements of $\mathfrak{su}(n)$ such that $X, X' \in (2\pi i\mathbb{Z})^n$. If ψ and ψ' denote the circle actions on $G_k(\mathbb{C}^n)$ defined by X and X' respectively. Then the equality*

$$[\psi] = [\psi'] \in [U(1), \text{Ham}(\mathcal{O})]_{gh}$$

implies that there exist a constant β and a bijective map $f : \mathcal{C}_k \rightarrow \mathcal{C}_k$, such that $\alpha'_{\mathfrak{d}} = \alpha_{f(\mathfrak{d})} + \beta$, for all $\mathfrak{d} \in \mathcal{C}_k$.

Remarks. When $k = 1$ the necessary condition stated in Theorem 12 for the equality $[\psi] = [\psi']$ is precisely the condition (c) of Theorem 11, since in this case the constant $\beta = a'_j - a_{f(j)}$ vanishes.

On the other hand the mentioned necessary condition given in Theorem 12 is independent of the symplectic structure. In Proposition 37 we give such a necessary condition when the orbit is diffeomorphic to the full flag manifold for $SU(n)$, which depends on the symplectic structure.

The paper is organized as follows. In Section 2 we introduce the cohomology class S_ψ and we prove Theorem 3 and Proposition 4.

Section 3 deals with the case when the manifold is the total space of the fibration (1.1); in particular Theorem 6 is proved. Subsection 3.1 is devoted to prove Theorem 7. The case $M = S^2 \times S^2$ is considered in Subsection 3.2.

In Section 4 we will concern with the relation $\sim$ between circle actions in the coadjoint orbits of a Lie group G , defined by elements of a Cartan subalgebra of G . We will prove the general Theorem 9, and Theorem 11 and Theorem 12 relative to circle actions on coadjoint orbits of $SU(n)$.

Acknowledgements. The relation $\sim$ has been introduced in [23]. I thank to a referee of paper [23] for suggesting me to study the relation between the space $[G, \text{Ham}(M)]_{gh}$ and the r -equivalence of G -actions introduced above. I thank the referee for his useful comments.

2. THE COHOMOLOGY CLASS S_ψ

Here we will denote by $\mathcal{H}$ the Hamiltonian group $\text{Ham}(M, \omega)$. From the universal principal bundle $E\mathcal{H} \rightarrow B\mathcal{H}$ of the group $\mathcal{H}$, we construct the universal bundle with fibre M

$$\begin{array}{ccc} M & \longrightarrow & M_{\mathcal{H}} := E\mathcal{H} \times_{\mathcal{H}} M \\ & & \downarrow \pi_{\mathcal{H}} \\ & & B\mathcal{H}. \end{array}$$

There exists a unique class $\mathbf{c} \in H^2(M_{\mathcal{H}}, \mathbb{R})$ called the coupling class (see [14]) satisfying:

- a) $\pi_{\mathcal{H}*}(\mathbf{c}^{n+1}) = 0$, where $\pi_{\mathcal{H}*}$ is the fibre integration.
- b) $\mathbf{c}$ extends the fiberwise class $[\omega]$.

Let G be a compact Lie group. We assume that M is equipped with the Hamiltonian G -action given by the group homomorphism $\psi : G \rightarrow \text{Ham}(M, \omega)$. This homomorphism induces a map $\Psi : BG \rightarrow B\mathcal{H}$, between the corresponding classifying spaces. The pullback $\Psi^{-1}(M_{\mathcal{H}})$ of $M_{\mathcal{H}}$ by Ψ is a bundle over BG which can be identified with $p : M_\psi = EG \times_G M \rightarrow BG$. Thus we have the following commutative diagram

$$\begin{array}{ccc} M_\psi & \xrightarrow{\hat{\Psi}} & M_{\mathcal{H}} \\ p \downarrow & & \downarrow \pi_{\mathcal{H}} \\ BG & \xrightarrow{\Psi} & B\mathcal{H}. \end{array}$$

By c_ψ we denote the pullback of $\mathbf{c}$ by $\hat{\Psi}$; that is, $c_\psi = \hat{\Psi}^*(\mathbf{c})$. The class c_ψ is the coupling class of the fibration $p : M_\psi \rightarrow BG$.

If ψ and ψ' are homotopic by means of a family of Hamiltonian G -actions, then the bundles $\Psi'^{-1}(M_{\mathcal{H}})$ and $\Psi^{-1}(M_{\mathcal{H}})$ are isomorphic and the isomorphism from M_ψ to $M_{\psi'}$ applies $c_{\psi'}$ to c_ψ .

Since $c_\psi \in H^2(M_\psi) = H_G^2(M)$ we can integrate e^{c_ψ} along the fibre and we obtain $S_\psi := p_*(e^{c_\psi})$ in the cohomology of $H(BG)$. *The above arguments prove Theorem 1.*

Strictly speaking e^{c_ψ} is not an equivariant de Rham cohomology class. For instance, when G is the n -torus T , e^{c_ψ} belongs to $\hat{H}_T(M)$ [3], where $\hat{H}_T(M)$ is

$$\hat{H}_T(M) = H_T(M) \otimes_{\mathbb{R}[u_1, \dots, u_n]} \mathbb{R}[[u_1, \dots, u_n]],$$

$\mathbb{R}[[u_1, \dots, u_n]]$ being the ring of formal power series in the variables $u_1, \dots, u_n$.

By the localization theorem [3] we have, in a suitable localization of $\hat{H}(BT) = \mathbb{R}[[u_1, \dots, u_n]]$,

$$(2.1) \quad S_\psi = \sum_F p_*^F \left(\frac{e^{c_\psi}}{E(N_F)} \right),$$

where F varies in the set of connected components of the fixed point set of the T -action, $p_*^F : \hat{H}_T(F) \rightarrow \hat{H}(BT)$ is the fibre integration on F , and $E(N_F)$ is the equivariant Euler class of the normal bundle N_F to F in M .

Let us assume that M is endowed with a Hamiltonian action of the torus $T = U(1)^n$. We say that a moment map $\mu : M \rightarrow \mathfrak{t}^*$ for the T -action is the *normalized* moment map if $\int_M \mu(X) \omega^n = 0$, for all $X \in \mathfrak{t}$. If Δ denotes the moment polytope of μ , that is $\Delta = \text{Im}(\mu)$, then

$$\int_M \mu \omega^n = \text{Cm}(\Delta) \int_M \omega^n,$$

where $\text{Cm}(\Delta)$ is the center of mass of Δ . It follows that $\text{Cm}(\Delta) = 0$, iff μ is normalized.

If μ is a normalized moment for the T -action ψ , then $\omega - \mu$ is a T -equivariant closed 2-form [13] which represents the coupling class c_ψ , because it satisfies the properties which determine the coupling class uniquely.

Let M be a toric manifold and ψ the corresponding effective action of T . We denote by Δ the respective moment polytope with center of mass at 0. By $u_1, \dots, u_n$ we denote the basis of $\mathfrak{t}^*$ dual of $e_1, \dots, e_n \in \mathfrak{t} = \mathbb{R}^n$. The image by moment map μ of the fixed points $q_1, \dots, q_s$ of the T -action ψ are the vertices of Δ [2], [11]. The equivariant Euler class $E(N_{q_i})$ of the normal bundle to q_i in M is

$$E(N_{q_i}) = \prod_{j=1}^n l_{ij},$$

where the l_{ij} are the weights of isotropy representation of T at the tangent space T_{q_i} (see [13]). So $E(N_{q_i})$ is a homogeneous degree n polynomial in the variables $u_1, \dots, u_n$.

On the other hand $\mu(q_i) = \sum_j \beta_{ij} u_j$, where $(\beta_{ij})_j$ are the coordinates of the vertex $\mu(q_i)$. Let f be the polynomial $\prod_i \prod_j l_{ij}$. Denoting by $[S_\psi]_f$ the class of S_ψ

in the localization $\hat{H}(BT)_f$ (see [3]) it follows from (2.1)

$$(2.2) \quad [S_\psi]_f = \sum_{i=1}^s \frac{e^{-\sum_j \beta_{ij} u_j}}{\prod_j l_{ij}} \in \hat{H}(BT)_f.$$

So we can write

$$(2.3) \quad [S_\psi]_f = \frac{A}{f} \in \hat{H}(BT)_f,$$

where A is a linear combination of elements of

$$\mathcal{E} := \{E_i := \exp(-\sum_j \beta_{ij} u_j)\}_i$$

with coefficients in the polynomial ring $\mathbb{R}[u_1, \dots, u_n]$.

Let ψ' be another effective action of T on M , and f' be the polynomial defined by the equivariant Euler classes of the normal bundles of the fixed points. So

$$(2.4) \quad [S_{\psi'}]_{f'} = \frac{C}{f'} \in \hat{H}(BT)_{f'}$$

In order to compare $[S_\psi]_f$ and $[S_{\psi'}]_{f'}$ we introduce the extension $\mathcal{A}$ of $\hat{H}(BT)$

$$\mathcal{A} := \mathbb{R}[[u_1, \dots, u_n]] \otimes_{\mathbb{R}[u_1, \dots, u_n]} \mathbb{R}(u_1, \dots, u_n),$$

where $\mathbb{R}(u_1, \dots, u_n)$ is the field of rational functions in the variables $u_1, \dots, u_n$.

We have the following obvious Lemma

Lemma 13. *Suppose that $S_\psi = S_{\psi'} \in \hat{H}(BT)$. If $[S_\psi]_f = \frac{\tilde{A}}{f^p} \in \hat{H}(BT)_f$ and $[S_{\psi'}]_{f'} = \frac{\tilde{C}}{(f')^q} \in \hat{H}(BT)_{f'}$, with $p, q \in \mathbb{Z}$, then there exists a rational function $g \in \mathbb{R}(u_1, \dots, u_n)$ such that $\tilde{A} = \tilde{C}g$ in $\mathcal{A}$.*

Proof. $g = f^p/(f')^q$ satisfies the required condition. $\square$

Let $\{\gamma_{ij} \mid i = 1, \dots, r; j = 1, \dots, m\}$ be a family of real numbers. We put

$$F_i(u_1, \dots, u_m) = \exp\left(\sum_j \gamma_{ij} u_j\right).$$

We will prove that the set of exponential functions $\{F_i\} \in \mathcal{A}$ is $\mathbb{R}(u_1, \dots, u_m)$ -independent.

Lemma 14. *If $\{Q_i(u_1, \dots, u_m)\}_{i=1, \dots, r}$ is a set of rational functions, such that $\sum_i Q_i F_i = 0$ and $F_i \neq F_k$, for $i \neq k$, then $Q_i = 0$ for $i = 1, \dots, r$.*

Proof. We define the linear functions $h_i(u_1, \dots, u_m) = \sum_j \gamma_{ij} u_j$. We write $|\gamma_i| = (\sum_j \gamma_{ij}^2)^{1/2}$. Let us assume that $|\gamma_1| \geq |\gamma_i|$ for $i = 1, \dots, r$. Then $|\text{grad}(h_i)| \leq |\text{grad}(h_1)|$ and moreover $\text{grad}(h_i) \neq \text{grad}(h_1)$, for $i \neq 1$ by the assumption. Hence the vector $v := (\gamma_{11}, \dots, \gamma_{1m}) \in \mathbb{R}^m$ defines a direction such that

$$(2.5) \quad \lim_{\lambda \rightarrow +\infty} \frac{F_i(\lambda v)}{F_1(\lambda v)} = 0,$$

for $i \neq 1$.

If Q_1 would be nonzero

$$(2.6) \quad 1 = -\sum_{i>1} \frac{Q_i}{Q_1} \frac{F_i}{F_1}.$$

If we take the limit along λv as $\lambda \rightarrow +\infty$ in (2.6) we arrive to a contradiction, by the exponential decay (2.5). Hence the initial linear combination reduces to $\sum_{i>1} Q_i F_i = 0$. By repeating the argument it follows $Q_i = 0$ for all i . $\square$

Sometimes we will use particular cases of Lemma 14; for example, the $\mathbb{R}[u, u^{-1}]$ -independence of a family of exponential functions $\{e^{\gamma_i u}\}_i$, with $\gamma_i \neq \gamma_j$ for $i \neq j$.

Proposition 15. *Let ψ and ψ' be toric actions on M and Δ and Δ' be the respective moment polytopes with center of mass at 0. Then $\Delta \neq \Delta'$ iff $S_\psi \neq S_{\psi'}$.*

Proof. Let us suppose that $\Delta \neq \Delta'$. According to (2.2), the element A in (2.3) is a linear combination $A = \sum_i g_i E_i$, with $0 \neq g_i \in \mathbb{R}[u_1, \dots, u_n]$. Similarly, C in (2.4) is a linear combination of elements of

$$\mathcal{E}' := \{E'_i := \exp(-\sum_j \beta'_{ij} u_j)\}_i.$$

As the moment polytopes are not equal, there is a vertex in Δ which does not belong to Δ' . So there is an element, say E_k , in $\mathcal{E}$ which is not in $\mathcal{E}'$.

If there were a rational function $g \in \mathbb{R}(u_1, \dots, u_n)$ such that $A = Cg$ in $\mathcal{A}$, then $0 = A - Cg$ is a linear combination in $\mathcal{A}$ with coefficients in $\mathbb{R}(u_1, \dots, u_n)$, in which E_k appears once with the coefficient $g_k \neq 0$. That is,

$$(2.7) \quad 0 = A - Cg = g_k E_k + \sum_{G_r \in \mathcal{E}''} \tilde{g}_r G_r,$$

where $\mathcal{E}'' = (\mathcal{E} \cup \mathcal{E}') \setminus \{E_k\}$. By the $\mathbb{R}(u_1, \dots, u_n)$ -linear independence of the set $\mathcal{E}'' \cup \{E_k\}$, it follows from (2.7) that $g_k = 0$. This contradiction shows that there is no a rational function g such that $A = Cg$. It follows from Lemma 13 that $S_\psi \neq S_{\psi'}$.

On the other hand, if $\Delta = \Delta'$, then trivially $\beta_{ij} = \beta'_{ij}$. Moreover the equality of the polytopes implies the equality of the weights of the respective isotropy representations at the fixed points [11]. Hence, by (2.2), $[S_\psi]_f = [S_{\psi'}]_{f'}$. As the polynomial f and f' are equal, it turns out that $S_\psi = S_{\psi'}$. $\square$

Theorem 16. *Let ψ and ψ' be toric actions on M . Let Δ and Δ' be the respective moment polytopes with center of mass at 0. If $\Delta \neq \Delta'$, then $[\psi] \neq [\psi'] \in [T, \text{Ham}(M)]_{\text{gh}}$.*

Proof. It is a consequence of Proposition 15 and Theorem 1. $\square$

Proposition 17. *If ψ and ψ' are r -equivalent Hamiltonian G -actions on M , then $[\psi] = [\psi'] \in [G, \text{Ham}(M)]_{\text{gh}}$.*

Proof. There exists $h \in \text{Ham}(M)$ such that $h \circ \psi_g \circ h^{-1} = \psi'_g$, for all g . Let h_s be a path in $\text{Ham}(M)$ from Id to h , then $\{h_s \circ \psi \circ h_s^{-1}\}_s$ defines a homotopy of G -actions between ψ and ψ' . $\square$

Proof of Theorem 3.

Properties (iv) and (i) are equivalent by Proposition 15.

(v) “implies” (iv) by Theorem 1.

(iii) $\implies$ (v) is Proposition 17.

If $h \in \text{Ham}(M)$ satisfies $\psi'_g = h \circ \psi_g \circ h^{-1}$, for all $g \in T$, then the corresponding normalized moment maps satisfy $\mu' = \mu \circ h$. So $\Delta = \text{im } \mu = \text{im } \mu' = \Delta'$; that is, (iii) implies (ii).

Since M is a toric manifold $\text{Ham}(M) = \text{Symp}_0(M)$; thus (ii) implies (iii). $\square$

Lemma 18. *Let ψ be a toric action on M and Δ its moment polytope with center of mass at 0. If v is an automorphism of T , then there exists $V \in GL(n, \mathbb{Z})$ such that $V(\Delta)$ is the moment polytope with center of mass at 0 associated to the action $\psi \circ v$.*

Proof. Let v^* be the automorphism of $\mathfrak{t}^*$ induced by v . If μ is the normalized moment map of ψ , then $v^* \circ \mu$ is the corresponding moment map of $\psi \circ v$. So $V = v^*$ in the identification $\mathfrak{t}^* = (\mathbb{R}^n)^*$. $\square$

Proof of Proposition 4.

(a) “implies” (b). It is consequence of Proposition 17.

(b) “implies” (d). By definition of $\sim_{rp}$, there exists an automorphism v of T such that $\psi' \sim \psi \circ v$. By the implication (v) $\implies$ (i) of Theorem 3, $\Delta' = \Delta(\psi \circ v)$. The implication follows from Lemma 18.

(c) “implies” (d). If ψ' and ψ are conjugate then $\xi \circ (\psi \circ v)(g) \circ \xi^{-1} = \psi'(g)$, with ξ a symplectomorphism. By the proof of Lemma 18, $\mu' = v^* \circ \mu \circ \xi$. So $\Delta' = v^*(\Delta)$.

(d) “implies” (c). Let us assume that $V(\Delta) = \Delta'$ with $V \in GL(n, \mathbb{Z})$. V induces a group automorphism of T , which is denoted by v , such that $v^* = V$. So the actions $\psi \circ v$ and ψ' have the same moment polytope. By Delzant’s Theorem (Th. 2.1, [5]) there is a symplectomorphism ξ such that $\xi \circ (\psi \circ v)(g) \circ \xi^{-1} = \psi'(g)$; that is, ψ and ψ' are conjugate. $\square$

Let us assume that M is a symplectic manifold equipped with a Hamiltonian G -action, where G is a Lie group. Let X be an element of $\mathfrak{g}$ of Type (s) which determine a circle action ψ^X on M , and let F be a $2s$ -dimensional submanifold of M , connected component of the fixed point set of the circle action ψ^X . The normal bundle N_F to F in M decomposes in direct sum of $n - s$ $U(1)$ -equivariant line bundles

$$N_F = \bigoplus_{j=1}^{n-s} N_j.$$

On N_k $U(1)$ acts with a weight $p_k u \neq 0$, where u we denote the vector in $\mathfrak{u}(1)^*$ dual of the standard basis of $\mathfrak{u}(1)$. Then the $U(1)$ -equivariant Euler class of N_F is

$$E(N_F) = \prod_{k=1}^{n-s} (c_1(N_k) + p_k u) = u^{n-s} \prod_k p_k \prod_k \left(1 + \frac{c_1(N_k)}{p_k u}\right).$$

So

$$(2.8) \quad (E(N_F))^{-1} = u^{s-n} \left(\alpha_0 + \frac{\alpha_1}{u} + \frac{\alpha_2}{u^2} + \dots \right),$$

α_r being a $2r$ -form on F and $\alpha_0 \neq 0$. That is, $(E(N_F))^{-1}$ is a Laurent polynomial of degree $s - n$ with coefficients even dimensional forms on F .

On the other hand, if f is a normalized Hamiltonian for the $U(1)$ -action ψ^X , then $\omega - uf$ is a $U(1)$ -equivariant closed 2-form [13] which represents the coupling class c_{ψ^X} [16]. Hence the contribution of F to S_{ψ^X} in (2.1) is

$$\frac{e^{au}}{u^{n-s}} \int_F e^{\omega} \left(\alpha_0 + \frac{\alpha_1}{u} + \dots \right),$$

where $-a$ is the value of f on F . That is,

$$\frac{e^{au}}{u^{n-s}} \left(A_0 u + A_1 + \cdots + \frac{A_{m-1}}{u^{m-2}} \right),$$

with $A_0 \neq 0$.

Let F' be another component of the fixed point set. As $\text{codim } F' \geq \text{codim } F \geq 2$, there is a curve σ in M which joins F and F' and such that it does not meet other fixed point. So the only stationary points of the function $f \circ \sigma$ are the end points; that is, $f \circ \sigma$ is strictly monotone, and thus $f(F) \neq f(F')$. Therefore the contribution of F' to S_{ψ^X} will be of the form $e^{a'u} Q'(u)$, with $a \neq a'$ and $Q'(u)$ a Laurent polynomial of degree $(1/2) \dim F' - n$. We note that the contributions of F and F' can not be grouped together in the form $e^{\tilde{a}u} \tilde{Q}(u)$. So we have the following Lemma

Lemma 19. *If X is of Type (s) and $F_1, \dots, F_c$ are the $2s$ -dimensional components of the fixed point set of the circle action ψ^X , then*

$$S_{\psi^X} = \sum_{i=1}^c e^{a_i u} Q_i(u) + \sum_r e^{b_r u} \hat{Q}_r(u),$$

with $a_i \neq a_j$ for $i \neq j$, $Q_i(u)$ a Laurent polynomial of degree $s - n$ and $\hat{Q}_r(u)$ a Laurent polynomial of degree less than $s - n$.

Proof of Theorem 2. If X and Y are of different Type, say Type of $X = s$ greater than Type of Y , then in the expression for S_{ψ^X} given in Lemma 19 appears an exponential function with coefficient a Laurent polynomial of degree $s - n$, but all the Laurent polynomials in S_{ψ^Y} have degree less than $s - n$. The Theorem follows from the aforesaid $\mathbb{R}[u, u^{-1}]$ -linear independence of the exponential functions $\{e^{au}\}$ together with Theorem 1. $\square$

Let M be a toric manifold. To avoid trivial cases we assume that $\dim M > 2$. By ψ we denote the corresponding Hamiltonian effective T -action and by μ a moment map. In terms of the basis $\{u_j\}$ we write $\mu = \sum_j \mu_j u_j$. Given $X = \sum_j k_j e_j$ in the integer lattice of $\mathfrak{t}$, then $\mu^X = \langle \mu, X \rangle = \sum_j k_j \mu_j$ is a moment map for the $U(1)$ -action ψ^X defined as the composition of ψ with $z \in U(1) \mapsto (z^{k_j}) \in T$.

The following Proposition gives equivalent conditions to the property of being of type 0.

Proposition 20. *Let X be a vector in the integer lattice of $\mathfrak{t}$, the following are equivalent.*

- (α) *The set of fixed points for the $U(1)$ -action ψ^X is equal to the set of fixed points of the T -action ψ .*
- (β) *$\langle v, X \rangle \neq 0$, for all v parallel to an edge of the moment polytope Δ .*
- (γ) *$X \notin \mathfrak{g}_a = \text{Lie}(G_a)$, for all a , with G_a proper isotropy subgroup of the T -action ψ .*

Proof.

(α) $\implies$ (β). Let e be an edge of moment polytope Δ . e is of the form $p + tv$, with t in an interval of $\mathbb{R}$, p a vertex of e and $v \in \mathfrak{t}^*$. If $\langle v, X \rangle = 0$, then μ^X takes on $\mu^{-1}(e)$ the value $\langle p + tv, X \rangle = \langle p, X \rangle$. Thus the infinitely many points of $\mu^{-1}(e)$ are fixed by ψ^X . But the fixed point set of the T -action is $\{q_1, \dots, q_s\}$, the set of inverse images by μ of the vertices of Δ .

$(\beta) \implies (\gamma)$. If $X \in \mathfrak{g}_b$, we may assume that $\mathfrak{g}_b$ is maximal; that is, G_b is the stabilizer group of the points of $N := \mu^{-1}(e)$, with e an edge of Δ . So the induced vector field X_M on M vanishes on the submanifold N , and $d\mu^X = 0$ on this set. The values of the function μ^X on N are of the form $\langle p + tv, X \rangle$, with t varying in an interval of $\mathbb{R}$, and where p is a vertex of e and v is parallel to e . It follows from the vanishing of $d\mu^X|_N$ that $\langle v, X \rangle = 0$.

$(\gamma) \implies (\alpha)$. If X satisfies (γ) then the fixed point set for the $U(1)$ -action ψ^X is $\{q_1, \dots, q_s\}$ (see Corollary 10.9.1 in [13]). $\square$

If the equivalent properties stated in Proposition 20 hold, the fixed point set for ψ^X is $\{q_1, \dots, q_s\}$. The $U(1)$ -equivariant Euler class $E(N_{q_i})$ is a degree n monomial in a variable u , that we denote by l_i . If the moment map μ is normalized, then μ^X is the normalized Hamiltonian of the S^1 -action ψ^X . Therefore

$$[S_{\psi^X}] = \sum_{i=1}^s \frac{e^{-\beta_i u}}{l_i} \in \hat{H}(BU(1))_u,$$

with $\beta_i = \langle P_i, X \rangle$, and P_i the vertex of Δ image of q_i by μ . The proof of Theorem 5 is similar to the one given for Proposition 15.

Proof of Theorem 5. We put $\mathcal{E} = \{E_i = e^{-\beta_i u}\}$ and $\mathcal{E}' = \{E'_i = e^{-\beta'_i u}\}$, with $\beta'_i = \langle \mu(q_i), Y \rangle$. We write $[S_{\psi^X}] = A/u^n$ and $[S_{\psi^Y}] = C/u^n$, where $A, C \in \mathbb{R}[[u]]$ are linear combinations

$$A = \sum_{i=1}^s a_i E_i, \quad C = \sum_{i=1}^s c_i E'_i; \quad a_i \neq 0 \neq c_i, \quad \text{for all } i.$$

By hypothesis, there is an E_k that does not belong to $\mathcal{E}'$. Then, by the $\mathbb{R}$ -linear independence of the exponential functions, $A \neq C$. Thus $S_{\psi^X} \neq S_{\psi^Y}$, and by Theorem 1 $[\psi^X] \neq [\psi^Y] \in [U(1), \text{Ham}(M)]_{gh}$. $\square$

Proposition 21. *Let X, Y be elements in the integer lattice of $\mathfrak{t}$ that satisfy the conditions of Proposition 20. If*

$$(2.9) \quad \{\lambda \langle P_i, X \rangle : i = 1, \dots, s\} \neq \{\langle P_i, Y \rangle : i = 1, \dots, s\},$$

for all $0 \neq \lambda \in \mathbb{Z}$, where the P_i 's are the vertices of the moment polytope with center at 0. Then ψ^X and ψ^Y are S^1 -actions $\sim_{rp}$ -inequivalent.

Proof. Suppose the Proposition were false. Then there were an automorphism v of $U(1)$ such that $\psi^X \circ v \sim \psi^Y$. But $\psi^X \circ v = \psi^Z$, with $Z = v_*(X)$; that is, $Z = \lambda X$, with $0 \neq \lambda \in \mathbb{Z}$. Since $\langle P_i, Z \rangle = \lambda \langle P_i, X \rangle$, it follows from (2.9) together with Theorem 5 that $[\psi^Z] \neq [\psi^Y]$; that is, $\psi^X \circ v$ is not $\sim$ -equivalent to ψ^Y . This contradicts our supposition. $\square$

3. CIRCLE ACTIONS IN $\mathbb{P}(L_1 \oplus \dots \oplus L_{n-1} \oplus \mathbb{C})$.

We denote by $e_1, \dots, e_n$ the standard basis of $\mathbb{R}^n$. Given $a_1, \dots, a_{n-1} \in \mathbb{Z}$, we define the following vectors of $\mathbb{R}^n$:

$$v_j = -e_j, \quad j = 1, \dots, n, \quad v_{n+1} = \sum_{j=1}^{n-1} e_j, \quad v_{n+2} = e_n - \sum_{i=1}^{n-1} a_i e_i.$$

Let $\{k_i\}_{i=1,\dots,n+2}$ be a family of real numbers with $k_j = 0$, for $j = 1, \dots, n$, $k_{n+1} = \sigma > 0$, $k_{n+2} = \tau > 0$ satisfying $\tau + a_i\sigma > 0$ for all i , then we consider the following Delzant polytope Δ in $(\mathbb{R}^n)^*$

$$\Delta = \bigcap_{i=1}^{n+2} \{x \in (\mathbb{R}^n)^* : \langle x, v_i \rangle \leq k_i\}.$$

When $n = 3$ the polytope Δ is represented in the Figure 1.




FIGURE 1. The polytope Δ when $n = 3$

The symplectic manifold M_Δ associated to Δ is

$$M_\Delta = \{(z_i) \in \mathbb{C}^{n+2} : \sum_j' |z_j|^2 = \sigma/\pi, -\sum_{j=1}^{n-1} a_j |z_j|^2 + |z_n|^2 + |z_{n+2}|^2 = \tau/\pi\} / \simeq,$$

in the sum $\sum_j'$ the index j runs in the set $\{1, \dots, n-1, n+1\}$. The equivalence relation $\simeq$ is defined by $(z'_j) \simeq (z_j)$ iff there exist $\epsilon, \lambda \in U(1)$ such that

$$\begin{cases} z'_j = \lambda \epsilon^{-a_j} z_j, & j = 1, \dots, n-1; \\ z'_k = \epsilon z_k, & k = n, n+2; \\ z'_{n+1} = \lambda z_{n+1}. \end{cases}$$

The action of $U(1)^n$ on M_Δ given by

$$(3.1) \quad (\lambda_j) \cdot [z_i] = [\lambda_1 z_1, \dots, \lambda_n z_n, z_{n+1}, z_{n+2}]$$

defines a structure of toric manifold on M_Δ . Moreover the map μ on M_Δ defined by $\mu([z_j]) = \pi(|z_1|^2, \dots, |z_n|^2)$ is the corresponding moment map with $\text{im } \mu = \Delta$.

The manifold M_Δ is the total space of the fibration $\mathbb{P}(L_1 \oplus \dots \oplus L_{n-1} \oplus \mathbb{C}) \rightarrow \mathbb{C}P^1$, where L_i is the holomorphic line bundle over $\mathbb{C}P^1$ with first Chern number $-a_i$.

Given $[z] \in M_\Delta$, we write $w = (z_n, z_{n+2})$. The element in $\mathbb{P}(L_1 \oplus \dots \oplus L_{n-1} \oplus \mathbb{C})$ which corresponds to $[z]$ is the following one in the fibre over $[w] = [z_n : z_{n+2}] \in \mathbb{C}P^1$

$$(3.2) \quad [(z_n)^{a_1} z_1 : (z_{n+2})^{a_1} z_1 : \dots : (z_n)^{a_{n-1}} z_{n-1} : (z_{n+2})^{a_{n-1}} z_{n-1} : z_{n+1}]$$

We will write $([w], [w^{a_j} z_j : z_{n+1}])$ for the element in fibre bundle associated to $[z] \in M_\Delta$.

An element $\mathbf{b} = (b_1, \dots, b_n)$ in the integer lattice $\mathbb{Z}^n$ determines a Hamiltonian circle action $\psi_{\mathbf{b}}$ on M_Δ through the action (3.1). In the notation introduced above

$$\psi_{\mathbf{b}}(t)[z] = ([w_t], [(w_t)^{a_j} z_j \epsilon_j : z_{n+1}]),$$

where $\epsilon_k = \exp(2\pi i b_k t)$ and $w_t = (z_n \epsilon_n, z_{n+2})$.

Proof of Theorem 6. Let $\mathbf{b}' = (b'_1, \dots, b'_n)$ with

$$b'_j = b_j + a_j b_n, \text{ for } j = 1, \dots, n-1, \text{ and } b'_n = -b_n,$$

then

$$\psi_{\mathbf{b}'}(t)[z] = ([w'_t], [(w'_t)^{a_j} z_j \epsilon'_j : z_{n+1}]),$$

with $w'_t = (z_n \epsilon'_n, z_{n+2})$ and $\epsilon'_k = \exp(2\pi i b'_k t)$. So

$$\psi_{\mathbf{b}'}(t)[z] = ([u_t], [(u_t)^{a_j} z_j \epsilon_j : z_{n+1}]),$$

where $u_t = (z_n, z_{n+2} \epsilon_n)$.

Let $\{\hat{n}_s : s \in [0, 1]\}$ be a continuous curve of unitary vectors in $\mathbb{R}^3$, with $\hat{n}_0 = e_3$ and $\hat{n}_1 = -e_3$. We denote by $\xi^s(t)$ the rotation of $S^2 = \mathbb{C}P^1$ around the axis defined by the unitary vector $\hat{n}_s$ and angle $2\pi b_n t$. Then $\xi^0(t)[z_n : z_{n+2}] = [w_t]$ and $\xi^1(t)[z_n : z_{n+2}] = [u_t]$. We write $[w_{s,t}] := \xi^s(t)[z_n : z_{n+2}]$.

For each $s \in [0, 1]$ we define

$$\psi^s(t)[z] = ([w_{s,t}], [(w_{s,t})^{a_j} z_j \epsilon_j : z_{n+1}]).$$

Thus $\psi^0 = \psi_{\mathbf{b}}$ and $\psi^1 = \psi_{\mathbf{b}'}$. Since $\xi^s(t + t') = \xi^s(t) \circ \xi^s(t')$, the family $\{\psi^s\}$ is a homotopy between $\psi_{\mathbf{b}}$ and $\psi_{\mathbf{b}'}$ consisting of circle actions. $\square$

3.1. Hirzebruch surfaces. Here we will consider the manifold $\mathbb{P}(L_1 \oplus \dots \oplus L_{n-1} \oplus \mathbb{C})$ when $n = 2$. In this case Δ is the trapezoid in $(\mathbb{R}^2)^*$ with vertices $P_1(0, 0)$, $P_2(0, \tau)$, $P_3(\sigma, 0)$, $P_4(\sigma, \tau + a\sigma)$ and M_Δ is a Hirzebruch surface $\mathbb{P}(L_k \oplus \mathbb{C})$, where L_k the holomorphic line bundle over $\mathbb{C}P^1$ with first Chern number $k := -a$. More explicitly

$$M_\Delta = \{(z_j) \in \mathbb{C}^4 : |z_1|^2 + |z_3|^2 = \sigma/\pi, -a|z_1|^2 + |z_2|^2 + |z_4|^2 = \tau/\pi\} / \simeq,$$

with

$$(z_1, z_2, z_3, z_4) \simeq (\epsilon_1 \epsilon_2^{-a} z_1, \epsilon_2 z_2, \epsilon_1 z_3, \epsilon_2 z_4),$$

ϵ_i being an element of $U(1)$.

The map $[z_j] \mapsto ([z_2 : z_4], [z_2^{-a} z_3 : z_4^{-a} z_3 : z_1])$ gives a representation of M as a submanifold of $\mathbb{C}P^1 \times \mathbb{C}P^2$.

An element $\mathbf{b}$ in the lattice $\mathbb{Z}^2$ determines a Hamiltonian circle action $\psi_{\mathbf{b}}$ on M_Δ . The following Proposition is a particular case of Theorem 6

Proposition 22. *If $\mathbf{b} = (b_1, b_2)$ in the integer lattice $\mathbb{Z}^2$ and $\mathbf{b}' = (b_1 - kb_2, -b_2)$, then $\psi_{\mathbf{b}} \sim \psi_{\mathbf{b}'}$.*

Now we assume that $k = -a \neq 0$. The case $k = 0$ will be considered in Subsection 3.2. In order to calculate $S_{\mathbf{b}} := S_{\psi_{\mathbf{b}}}$ it is necessary to take into account the Type of $\mathbf{b}$. The element $\mathbf{b} \neq 0$ is of Type(0) when $\langle l, \mathbf{b} \rangle \neq 0$, for all edge l of Δ . And it is of Type (1) if there is an edge f such that $\langle f, \mathbf{b} \rangle = 0$.

Values of S_- on Type(0)

If $\mathbf{b} = (b_1, b_2) \in \mathbb{Z}^2$ is of type 0, then $b_1 \neq 0 \neq b_2 \neq 0 \neq b_0 := (b_1 + ab_2)$, and the fixed points of the S^1 -action $\psi_{\mathbf{b}}$ are the inverse images by the moment map μ of the vertices of Δ . If P is a vertex, the $U(1)$ -equivariant Euler class of the normal bundle of $\mu^{-1}(P)$ in M is the product of the weights of the corresponding isotropy representation. On the other hand, the edges meeting at P can be expressed as $\{P + t\rho_i : t \in [0, c_i]\}$, where $\rho_i \in (\mathbb{Z}^2)^*$ and ρ_1, ρ_2 is a basis of $(\mathbb{Z}^2)^*$. The aforementioned weights are precisely the numbers $\langle \rho_i, \mathbf{b} \rangle$, with $i = 1, 2$.

The normalized Hamiltonian function for the S^1 -action $\psi_{\mathbf{b}}$ is $\langle \mu, \mathbf{b} \rangle - \langle \text{Cm}(\Delta), \mathbf{b} \rangle$, where $\text{Cm}(\Delta)$ is the center of mass of Δ . Hence

$$(3.3) \quad S_{\mathbf{b}} = \frac{e^{qu}}{u^2} \sum_{j=1}^4 \frac{e^{-\langle P_j, \mathbf{b} \rangle u}}{\prod_{i=1}^2 \langle \rho_{ji}, \mathbf{b} \rangle},$$

with $\{\rho_{ji}\}_{i=1,2}$ is the basis of $(\mathbb{Z}^2)^*$ formed with the aforementioned vectors which give the directions of the edges meeting at P_j , and $q := \langle \text{Cm}(\Delta), \mathbf{b} \rangle$.

On $U_1 := \{[z] \in M : z_3 \neq 0 \neq z_4\}$ we can consider the complex coordinates $w = z_2/z_4$ and $w' = z_1/(z_3 z_4^k)$. Then

$$\psi_{\mathbf{b}}(w, w') = (e^{2\pi i b_2 t} w, e^{2\pi i b_1 t} w').$$

The point $\mu^{-1}(P_1)$ has coordinates $w = 0 = w'$, so the isotropy representation at this point has weights b_1 and b_2 ; in fact these values are the products by $\mathbf{b}$ of the direction vectors of edges meeting at P_1 . The remaining terms in (3.3) can be obtained in a similar way, or directly from the vectors ρ_i . So (3.3) can be written

$$(3.4) \quad u^2 S_{\mathbf{b}} = e^{qu} (A - B e^{-\tau b_2 u} - A e^{-\sigma b_1 u} + B e^{-(\tau b_2 + \sigma b_0)u})$$

where $A := (b_1 b_2)^{-1}$ and $B := (b_0 b_2)^{-1}$. Note that $A \neq B$; otherwise b_0 would be equal to b_1 , but this is impossible in Type(0).

The expression (3.4) will be written

$$(3.5) \quad u^2 S_{\mathbf{b}} = A e^{\alpha_1 u} - B e^{\alpha_2 u} - A e^{\alpha_3 u} + B e^{\alpha_4 u},$$

where $\alpha_1 = q$, $\alpha_2 = q - \tau b_2$, $\alpha_3 = q - \sigma b_1$, $\alpha_4 = q - (\tau b_2 + \sigma b_0)$. If $\alpha_2 \neq \alpha_3$ and $\alpha_1 \neq \alpha_4$, there are four different exponential functions in (3.5). By contrast, if $\alpha_2 = \alpha_3$ and $\alpha_1 = \alpha_4$ there are only two different exponentials in (3.5). Therefore we will distinguish three subtypes in Type(0).

$$\text{Type}(0) \begin{cases} \mathbf{b} \in \text{Subtype}(0)\alpha & \text{iff } \alpha_2 \neq \alpha_3, \alpha_1 \neq \alpha_4 \\ \mathbf{b} \in \text{Subtype}(0)\beta & \text{iff either } \alpha_2 = \alpha_3, \alpha_1 \neq \alpha_4, \text{ or } \alpha_2 \neq \alpha_3, \alpha_1 = \alpha_4 \\ \mathbf{b} \in \text{Subtype}(0)\gamma & \text{iff } \alpha_2 = \alpha_3, \alpha_1 = \alpha_4 \end{cases}$$

It follows from the Lemma 14 the following Proposition

Proposition 23. *If $\mathbf{b}$ and $\mathbf{b}'$ are in Type(0) and belong to different subtype, then $S_{\mathbf{b}} \neq S_{\mathbf{b}'}$.*

Given $\mathbf{b}, \mathbf{b}' \in \text{Subtype}(0)\alpha$, if $S_{\mathbf{b}} = S_{\mathbf{b}'}$, then by Lemma 14 it follows from (3.5)

$$(3.6) \quad \{A e^{\alpha_1 u}, -B e^{\alpha_2 u}, -A e^{\alpha_3 u}, B e^{\alpha_4 u}\} = \{A' e^{\alpha'_1 u}, -B' e^{\alpha'_2 u}, -A' e^{\alpha'_3 u}, B' e^{\alpha'_4 u}\}.$$

This set equality gives, in principle, the following eight possibilities

$$\mathbf{b}, \mathbf{b}' \in \text{Subtype}(0)\alpha \begin{cases} (i) A = A' & \begin{cases} (a) B = B' \\ (b) B = -B' \end{cases} \\ (ii) A = B' & \begin{cases} (a) B = A' \\ (b) B = -A' \end{cases} \\ (iii) A = -A' & \begin{cases} (a) B = B' \\ (b) B = -B' \end{cases} \\ (iv) A = -B' & \begin{cases} (a) B = A' \\ (b) B = -A' \end{cases} \end{cases}$$

Where the corresponding exponential function which multiplies each constant has been deleted in the equalities; for example, we have written $B = -A'$ instead of $Be^{\alpha_4 u} = -A'e^{\alpha_3 u}$.

Proposition 24. *If $\mathbf{b}$ and $\mathbf{b}'$ are in Subtype(0) α and $S_{\mathbf{b}} = S_{\mathbf{b}'}$, then $\mathbf{b} \equiv \mathbf{b}'$.*

Proof. In the case (i)(a), $A = A'$, $B = B'$, $\alpha_1 = \alpha'_1$, $\alpha_4 = \alpha'_4$. And by (3.6) $-Be^{\alpha_2 u} = -B'e^{\alpha'_2 u}$, $-Ae^{\alpha_3 u} = -A'e^{\alpha'_3 u}$. From $\alpha_j = \alpha'_j$ for all j , we deduce $\mathbf{b}' = \mathbf{b}$.

In the case (i)(b), $-Ae^{\alpha_3 u} = -A'e^{\alpha'_3 u}$. So $\alpha_3 = \alpha'_3$ and of course $\alpha_1 = \alpha'_1$. Thus $b_1 = b'_1$. From $A = A'$ it follows $b'_2 = b_2$; that is, $\mathbf{b} = \mathbf{b}'$. But this is contradictory with $B = -B'$.

It is also easy to check that the cases (ii)(a), (ii)(b), (iii)(a), (iii)(b) and (iv)(a) do not occur. In the case (iv)(b) it is straightforward to deduce $b'_1 = b_1 - kb_2$ and $b'_2 = -b_2$. $\square$

Next we study the map S_- on Subtype(0) β . If $\mathbf{b}$ belongs to Subtype(0) β , then either

$$u^2 S_{\mathbf{b}} = Ae^{\alpha_1 u} - (A + B)e^{\alpha_2 u} + Be^{\alpha_4 u}, \text{ when } \alpha_2 = \alpha_3$$

or

$$u^2 S_{\mathbf{b}} = (A + B)e^{\alpha_1 u} - Ae^{\alpha_2 u} - Be^{\alpha_3 u}, \text{ when } \alpha_1 = \alpha_4.$$

We will consider the partition $\mathcal{P}$ of Subtype(0) β formed by two subsets: The one consisting of all elements for which $\alpha_2 = \alpha_3$, and the other one its complement subset. It is not hard to prove the following Proposition

Proposition 25. *Given $\mathbf{b}, \mathbf{b}' \in \text{Subtype}(0)\beta$ with $S_{\mathbf{b}} = S_{\mathbf{b}'}$, if they belong to the same subset of the partition $\mathcal{P}$, then $\mathbf{b}' = \mathbf{b}$. If they belong to different subsets, then $\mathbf{b}' \equiv \mathbf{b}$.*

It is easy to prove

Proposition 26. *The map S_- is injective on Subtype(0) γ .*

Propositions 24, 25 and 26 can be put together

Proposition 27. *If $\mathbf{b}, \mathbf{b}' \in \text{Type}(0)$ and $S_{\mathbf{b}} = S_{\mathbf{b}'}$, then $\mathbf{b} \equiv \mathbf{b}'$.*

Values of S_- on Type(1)

If $\mathbf{b}$ belongs to Type(1) one of b_j 's vanishes. We will distinguish three possibilities: $b_2 = 0$, $b_1 = 0$ and $\mathbf{b} = (kb_2, b_2)$, that is $b_0 = 0$.

We first consider the case $b_2 = 0$. Now $\psi_{\mathbf{b}}[z] = [e^{2\pi i b_1 t} z_1, z_2, z_3, z_4]$. So the fixed point set is $\{[z] \in M : z_1 z_3 = 0\}$. On $U_2 = \{[z] \in M : z_2 \neq 0 \neq z_3\}$ we consider the complex coordinates $v = z_4/z_2$ and $v' = z_1/(z_3 z_2^k)$. So $\frac{\partial}{\partial v'}$ is a section of the normal bundle N_Q to $Q = \{[z] : z_1 = 0\}$ on $Q \cap U_2$. On the other hand, $\frac{\partial}{\partial w'}$ is a section of N_Q on $Q \cap U_1$. As

$$\frac{\partial}{\partial w'} = \left(\frac{z_4}{z_2}\right)^k \frac{\partial}{\partial v'},$$

identifying Q with $\mathbb{CP}^1$ it turns out that N_Q is $\mathcal{O}(k)$.

Since the coordinate w' is transformed by $\psi_{\mathbf{b}}(t)$ in $\exp(2\pi i b_1 t)w'$, the $U(1)$ -equivariant Euler class $E(N_Q)$ of the normal bundle to N_Q is $c_1(\mathcal{O}(k)) + b_1 u$.

Analogously, if we denote by $Q' := \{[z] : z_3 = 0\}$, then $E(N_{Q'}) = c_1(\mathcal{O}(-k)) - b_1 u$. As $\omega(Q) = \tau$ and $\omega(Q') = \tau - k\sigma =: \lambda$, it follows from the localization formula

$$(3.7) \quad S_{\mathbf{b}} = e^{qu} \left(\frac{1}{b_1 u} \left(\tau - \frac{k}{b_1 u} \right) - \frac{e^{-\sigma b_1 u}}{b_1 u} \left(\lambda - \frac{k}{b_1 u} \right) \right),$$

where $q := \langle \text{Cm}(\Delta), \mathbf{b} \rangle$.

Next we consider the case when $\mathbf{b} = (0, b_2)$, with $b_2 \neq 0$. Now the fixed point set is $\{[z] : z_2 = 0\} \cup \{P_2, P_4\}$. The vector field $\frac{\partial}{\partial w}$ is a global section to the normal bundle to $\{[z] : z_2 = 0\}$, so its $U(1)$ -equivariant Euler class is $b_2 u$. The equivariant Euler classes of the normal bundles N_{P_2} and N_{P_4} are $k(b_2 u)^2$ and $-k(b_2 u)^2$ respectively. As $\omega(\{[z] : z_2 = 0\}) = \sigma$, it follows

$$(3.8) \quad S_{\mathbf{b}} = e^{qu} \left(\frac{e^{-\tau b_2 u}}{k(b_2 u)^2} - \frac{e^{-\lambda b_2 u}}{k(b_2 u)^2} + \frac{\sigma}{b_2 u} \right).$$

Finally, let us suppose that $\mathbf{b} = (kb_2, b_2)$. The corresponding fixed point set is $\{[z] : z_4 = 0\} \cup \{P_1, P_3\}$. The equivariant Euler class of the normal bundles to $\{[z] : z_4 = 0\}$, P_1 and P_3 are $-b_2 u$, $k(b_2 u)^2$ and $-k(b_2 u)^2$ respectively. So

$$(3.9) \quad S_{\mathbf{b}} = e^{qu} \left(\frac{1}{k(b_2 u)^2} - \frac{e^{-\sigma k b_2 u}}{k(b_2 u)^2} - \frac{\sigma e^{-\tau b_2 u}}{b_2 u} \right).$$

To compare the expressions $S_{\mathbf{b}}$ for different elements $\mathbf{b}$, we classify the elements of $\text{Type}(0)$ according to the following scheme of subtypes

$$\text{Type}(1) \begin{cases} \mathbf{b} \in \text{Subtype}(1)\alpha & \text{iff } b_2 = 0 \\ \mathbf{b} \in \text{Subtype}(1)\beta & \text{iff } b_2 \neq 0 \end{cases}$$

From (3.7), (3.8) and (3.9) together with Lemma 14 it follows the following Proposition

Proposition 28. *If $\mathbf{b}, \mathbf{b}'$ belong to $\text{Type}(1)$ but to different subtype, then $S_{\mathbf{b}} \neq S_{\mathbf{b}'}$.*

It follows from Theorem 1 together with Theorem 2, Proposition 28 and Proposition 23 the following Theorem

Theorem 29. *Let M_{Δ} be a Hirzebruch surface non diffeomorphic to $S^2 \times S^2$. Let $\mathbf{b}$ and $\mathbf{b}'$ be nonzero elements of $\mathbb{Z}^2$. If they belong to different subtype then $\psi_{\mathbf{b}} \not\sim \psi_{\mathbf{b}'}$.*

Next we study whether the map $\mathbf{b} \mapsto S_{\mathbf{b}}$ is injective restricted to each of the different subtypes of $\text{Type}(1)$.

If $\mathbf{b} = (b_1, 0)$ belongs to $\text{Subtype}(1)\alpha$, in the expression of $u^2 S_{\mathbf{b}}$ obtained from (3.7) there are two different exponentials

$$u^2 S_{\mathbf{b}} = A(u) e^{qu} + B(u) e^{(q - \sigma b_1)u},$$

where $A(u)$ and $B(u)$ are the following polynomials $A(u) = (b_1)^{-1} \tau u - k(b_1)^{-2}$, $B(u) = -(b_1)^{-1} \lambda u + k(b_1)^{-2}$. If $\mathbf{b}'$ belongs to $\text{Subtype}(1)\alpha$ and $S_{\mathbf{b}} = S_{\mathbf{b}'}$ then either $A(u) = A'(u)$ and $B(u) = B'(u)$ or $A = B'$ and $B = A'$. In the first case $b'_1 = b_1$. In the second one implies $-(b_1)^2 = (b'_1)^2$. Hence we have the following Proposition

Proposition 30. *S_- is injective on $\text{Subtype}(1)\alpha$.*

In Subtype(1) β one can distinguish the case in which $\mathbf{b}$ is of the form $(0, c)$ and when it is of the form (kc, c) . As in the preceding Proposition, if $\mathbf{b} = (0, b_2)$, $\mathbf{b}' = (0, b'_2)$ and $S_{\mathbf{b}} = S_{\mathbf{b}'}$, then $\mathbf{b} = \mathbf{b}'$. Analogously when $\mathbf{b} = (kc, c)$ and $\mathbf{b}' = (kc', c')$.

Finally, given $\mathbf{b} = (0, b_2)$ and $\mathbf{b}' = (b'_1, b'_2) = (kc, c)$, in the expressions for $u^2 S_{\mathbf{b}}$ and $u^2 S_{\mathbf{b}'}$ obtained from (3.8) and (3.9) there are three different exponentials

$$\begin{aligned} u^2 S_{\mathbf{b}} &= Ae^{(q-\tau b_2)u} + Be^{(q-\lambda b_2)u} + C(u)e^{qu}, \\ u^2 S_{\mathbf{b}'} &= A'e^{q'u} + B'e^{(q'-\sigma k b'_2)u} + C'(u)e^{(q'-\tau b'_2)u}, \end{aligned}$$

with $C(u) = b_2^{-1}\sigma$ and $C'(u) = -(b'_2)^{-1}\sigma$. The equality $S_{\mathbf{b}} = S_{\mathbf{b}'}$ implies $C(u) = C'(u)$; that is $b'_2 = -b_2$. In other words $\mathbf{b} \equiv \mathbf{b}'$. We have proved the Proposition

Proposition 31. *If $\mathbf{b}$ and $\mathbf{b}'$ belong to Subtype(1) β with $S_{\mathbf{b}} = S_{\mathbf{b}'}$ then $\mathbf{b} \equiv \mathbf{b}'$.*

The preceding Propositions can be summarize in the following one

Proposition 32. *If $\mathbf{b}$ and $\mathbf{b}'$ are elements of type 1 with $S_{\mathbf{b}} = S_{\mathbf{b}'}$ then $\mathbf{b} \equiv \mathbf{b}'$.*

Proof of Theorem 7. Theorem 7 is consequence of Proposition 32, Proposition 27, Proposition 22 and Theorem 1. $\square$

3.2. The manifold $S^2 \times S^2$. Here we study the relation $\psi_{\mathbf{b}} \sim \psi_{\mathbf{b}'}$ when M is the symplectic manifold associated to the polytope Δ in the case $a = 0$. That is, M is diffeomorphic to $S^2 \times S^2$. Now $b_0 = b_1$ and $\text{Cm}(\Delta) = (\sigma/2, \tau/2)$, so the corresponding discussion is simpler than the one of case $a \neq 0$.

If $b_1 \neq 0 \neq b_2$, the expression (3.4) (which is also valid when $k = 0$) reduces to

$$(3.10) \quad S_{\mathbf{b}} = \frac{4}{b_1 b_2 u^2} \left(\sinh\left(\frac{\sigma b_1 u}{2}\right) \sinh\left(\frac{\tau b_2 u}{2}\right) \right)$$

When $\mathbf{b} = (b_1, 0)$ the expression for $S_{\mathbf{b}}$ can be obtained from (3.7)

$$(3.11) \quad S_{\mathbf{b}} = \frac{2\tau}{b_1 u} \sinh\left(\frac{\sigma b_1 u}{2}\right)$$

Analogously, if $\mathbf{b} = (0, b_2)$, then

$$(3.12) \quad S_{\mathbf{b}} = \frac{2\sigma}{b_2 u} \sinh\left(\frac{\tau b_2 u}{2}\right)$$

Using Lemma 14 it is straightforward to deduce from (3.10) the following Proposition

Proposition 33. *If $b_1 b_2 \neq 0 \neq b'_1 b'_2$, then $S_{\mathbf{b}} = S_{\mathbf{b}'}$ iff one of the following statements are true:*

- (i) $|b'_1| = |b_1|$ and $|b'_2| = |b_2|$.
- (ii) $\sigma b_1 = \pm \tau b'_2$ and $\tau b_2 = \pm \sigma b'_1$.
- (iii) $\sigma b_1 = \pm \tau b'_2$ and $\tau b_2 = \mp \sigma b'_1$.

Proof of Theorem 8. In this case

$$\psi_{\mathbf{b}}(t)[z] = ([e^{2\pi i b_2 t} z_2 : z_4], [z_3 : e^{2\pi i b_1 t} z_1]),$$

where $[z] = ([z_2 : z_4], [z_3 : z_1]) \in S^2 \times S^2$. On the other hand, if $\sigma/\tau \notin \mathbb{Q}$, the possibilities (ii) and (iii) in the preceding Proposition do not occur. If, for example $b'_1 = -b_1$ and $b'_2 = b_2$, let ξ_t^s be the rotation of angle $2\pi b_1 t$ around the vector $\hat{n}_s$ introduced in the Proof of Theorem 6. We set

$$\psi_t^s[z] = ([e^{2\pi i b_2 t} z_2 : z_4], \xi_t^s[z_3 : z_1]).$$

Then ψ^s is a homotopy between $\psi_{\mathbf{b}}$ and $\psi_{\mathbf{b}'}$ consisting of $U(1)$ -actions. That is, $|b'_1| = |b_1|$ and $|b'_2| = |b_2|$ implies $\psi_{\mathbf{b}} \sim \psi_{\mathbf{b}'}$. So Theorem 8 follows from Proposition 33 together with Theorem 1. $\square$

Remark. In general $S_{\mathbf{b}} = S_{\mathbf{b}'}$ does not imply $[\psi_{\mathbf{b}}] = [\psi_{\mathbf{b}'}] \in [U(1), \text{Ham}(M)]$. For instance, if $\tau = \sigma$, $\mathbf{b} = (1, 0)$ and $\mathbf{b}' = (0, 1)$, it follows from (3.11) and (3.12) that $S_{\mathbf{b}} = S_{\mathbf{b}'}$. Gromov proved that $\text{Ham}(S^2 \times S^2)$ deformation retracts to the group $SO(3) \times SO(3)$ [8]. So $\pi_1(\text{Ham}(S^2 \times S^2)) = \mathbb{Z}_2 \times \mathbb{Z}_2$ and $\psi_{\mathbf{b}}$ and $\psi_{\mathbf{b}'}$ are the generators of the respective groups in the product. Hence $\psi_{\mathbf{b}}$ and $\psi_{\mathbf{b}'}$ are not homotopic, and a fortiori $\psi_{\mathbf{b}} \not\sim \psi_{\mathbf{b}'}$.

4. S^1 -ACTIONS IN COADJOINT ORBITS.

Let G be a compact connected Lie group and $\eta \in \mathfrak{g}^*$. We consider the coadjoint orbit $\mathcal{O}$ of the element η , endowed with the standard symplectic structure ω (see [17]). A moment map $\mu_G : \mathcal{O} \rightarrow \mathfrak{g}^*$ for the action of G on $\mathcal{O}$ is the opposite of the inclusion. The manifold $\mathcal{O}$ can be identified with the quotient G/G_η , where G_η is the subgroup of isotropy of η . Let H be a Cartan subgroup of G contained in G_η ([12], page 166). We have the decomposition of $\mathfrak{g}_{\mathbb{C}}$ as direct sum of root spaces

$$\mathfrak{g}_{\mathbb{C}} = \mathfrak{h}_{\mathbb{C}} \oplus \bigoplus_{\alpha \in \mathcal{R}} \mathfrak{g}_{\alpha},$$

where $\mathcal{R}$ is the set of roots determined by H . We will denote by $\check{\alpha} \in \mathfrak{h}$ the coroot associated to $\alpha \in \mathcal{R}$ [7]. Let $\mathfrak{p}$ be the parabolic subalgebra

$$\mathfrak{p} = \mathfrak{h}_{\mathbb{C}} \oplus \bigoplus_{\eta(i\check{\alpha}) \leq 0} \mathfrak{g}_{\alpha}.$$

If P is the parabolic subgroup of $G_{\mathbb{C}}$ generated by $\mathfrak{p}$, then $G_{\mathbb{C}}/P$ is a complexification of $\mathcal{O}$ and

$$(4.1) \quad T_{\eta}^{1,0} \mathcal{O} = \bigoplus_{\alpha \in \mathcal{R}^+} \mathfrak{g}_{\alpha},$$

where $\mathcal{R}^+ = \{\alpha \in \mathcal{R} : \langle \eta, i\check{\alpha} \rangle > 0\}$.

Given $X \in \mathfrak{g}$ we consider the map $h := \langle \mu_G, X \rangle : \mathcal{O} \rightarrow \mathbb{R}$, and define the constant $\kappa \in \mathbb{R}$ by the relation

$$\kappa \int_{\mathcal{O}} \omega^n = \int_{\mathcal{O}} h \omega^n.$$

The function $h - \kappa$ is the normalized Hamiltonian function for the isotopy in $\mathcal{O}$ generated by X through the G -action.

If X belongs to the integer lattice of $\mathfrak{h}$, then X determines a $U(1)$ -action ψ on $\mathcal{O}$. By u we denote a coordinate in $\mathfrak{u}(1)$, then closed $U(1)$ -equivariant 2-form $\omega - (\mu - \kappa)u$ is a representative of the coupling class c_{ψ} [16]. Thus

$$S_{\psi} = e^{\kappa u} \int_{\mathcal{O}} e^{-\mu u + \omega} \in \hat{H}(BU(1)).$$

In order to apply the localization formula we need to determine the fixed point set of the S^1 -action ψ . Let us assume that X is a regular element [6] of the lattice of $\mathfrak{h}$. We denote by W_{η} the stabilizer subgroup of η in the Weyl group W of (G, H) . It is easy to see that the fixed point set of the action ψ is $\{w \cdot \eta : w \in W/W_{\eta}\}$ (see [4] page 231).

Given $w \in W$, the $U(1)$ -equivariant Euler class of the normal bundle to the point $w \cdot \eta$ in $\mathcal{O}$ can be calculated as follows: From (4.1) it follows

$$T_{w \cdot \eta}^{1,0} \mathcal{O} = \bigoplus_{\alpha \in \mathcal{R}^+} \mathfrak{g}_{w \cdot \alpha}.$$

Moreover $[Y, Z] = (w \cdot \alpha)(Y)Z$ for any $Z \in \mathfrak{g}_{w \cdot \alpha}$ and all $Y \in \mathfrak{h}$. So the representation of $U(1)$ on $\mathfrak{g}_{w \cdot \alpha}$ induced by ψ therefore has the weight $-i(w \cdot \alpha) \in \mathfrak{u}(1)^*$, and the weights of the isotropy representation at the point $w \cdot \eta \in \mathcal{O}$ are $-i(w \cdot \alpha)$, with $\alpha \in \mathcal{R}^+$. That is, the $U(1)$ -equivariant Euler class satisfies

$$(4.2) \quad E(N_{w \cdot \eta})(Y) = \prod_{\alpha \in \mathcal{R}^+} -i(w \cdot \alpha)(Y),$$

for any $Y = tX$ with $t \in \mathbb{R}$. Thus $E(N_{w \cdot \eta})$ is a monomial in u

$$(4.3) \quad E(N_{w \cdot \eta}) = \prod_{\alpha \in \mathcal{R}^+} A_{\alpha}^w u \in H(BU(1)),$$

with A_{α}^w constant.

Consequently, if X is a regular element of the integer lattice of $\mathfrak{h}$, in the localization $\hat{H}(BU(1))_u$ one has

$$(4.4) \quad S_{\psi} = e^{\kappa u} \sum_{w \in W/W_{\eta}} \frac{e^{\eta(w^{-1} \cdot X)u}}{\prod_{\alpha \in \mathcal{R}^+} A_{\alpha}^w u},$$

since $\mu(w \cdot \eta) = -\langle w \cdot \eta, X \rangle = -\eta(w^{-1} \cdot X)$.

Theorem 34. *Given X, X' regular elements of the integer lattice of $\mathfrak{h}$, let ψ and ψ' be the circle actions on $\mathcal{O}$ generated by X and X' , respectively. If there is no a translation in $\mathbb{R}$ which applies the set of real numbers $\{\eta(w \cdot X) : w \in W/W_{\eta}\}$ in $\{\eta(w \cdot X') : w \in W/W_{\eta}\}$, then ψ and ψ' are not homotopically equivalent by means of a homotopy consisting of circle actions*

Proof. The class $S_{\psi'}$ is obtained from (4.4) by substituting κ, X and A_{α}^w by the corresponding κ', X' and $A_{\alpha}'^w$. By the hypothesis the sets of functions

$$\mathcal{E} := \{e^{(\kappa + \eta(w^{-1} \cdot X))u} \mid w \in W/W_{\eta}\} \text{ and } \mathcal{E}' := \{e^{(\kappa' + \eta(w^{-1} \cdot X'))u} \mid w \in W/W_{\eta}\}$$

are distinct. Hence $\mathcal{E}$ is not contained in the $\mathbb{R}[u]$ -submodule of $\mathbb{R}[[u]]$ generated by $\mathcal{E}'$. Reasoning as in the proof of Theorem 5 we conclude that $S_{\psi} \neq S_{\psi'}$, and the theorem follows from Theorem 1. $\square$

Proposition 35. *Let ψ and ψ' be the circle actions on the coadjoint orbit $\mathcal{O}$ generated by the regular elements X and X' of the integer lattice of $\mathfrak{h}$, respectively. If there is no an affine transformation*

$$B : x \in \mathbb{R} \rightarrow \lambda x + a \in \mathbb{R}, \text{ with } 0 \neq \lambda \in \mathbb{Z} \text{ and } a \in \mathbb{R},$$

such that

$$B(\{\eta(w \cdot X) : w \in W/W_{\eta}\}) = \{\eta(w \cdot X') : w \in W/W_{\eta}\},$$

then ψ and ψ' are $\sim_{rp}$ -inequivalent.

Proof. If ψ and ψ' were $\sim_{rp}$ -equivalent, there would be an automorphism v of $U(1)$ with $\psi' \sim \psi \circ v$. Moreover $\psi \circ v$ is the S^1 -action generated by $Z = cX$, with c a nonzero integer. By Theorem 34 there must be a translation T_b in $\mathbb{R}$ which applies $\{\eta(w \cdot Z) = c\eta(w \cdot X) : w \in W/W_\eta\}$ in $\{\eta(w \cdot X') : w \in W/W_\eta\}$. Hence the affine transformation $B : x \in \mathbb{R} \mapsto cx + b \in \mathbb{R}$ contradicts the hypothesis of the Proposition. $\square$

Proof of Theorem 9. We consider the coadjoint orbit $\mathcal{O} = G/Q$ of G , where Q is a subgroup of G which contains a maximal torus H of G . The element X in the integer lattice of $\mathfrak{h}$ generates the circle action

$$\psi_t : gQ \in G/Q \mapsto (e^{tX}g)Q \in G/Q.$$

We denote by ψ' the S^1 -action defined by X' . Let g_1 be an element of the normalizer $N(H)$ of H in G , such that $g_1H \in N(H)/H = W$ corresponds to the element w . And let $g : s \in [0, 1] \mapsto g(s) \in G$ be a curve with $g(0) = 1$ and $g(1) = g_1$. We write $X^s := \text{Ad}_g(s)X$, then $X^1 = w(X) = X'$. For $s, t \in [0, 1]$ we put

$$\psi_t^s : gQ \in G/Q \mapsto (e^{tX^s}g)Q \in G/Q.$$

As $e^{X^s} = g(s)e^X(g(s))^{-1} = 1$, for each s the family $\{\psi_t^s\}_t$ is a circle action on the orbit $\mathcal{O} = G/Q$. Moreover

$$\psi_t^1(gQ) = (e^{tX^1}g)Q = (e^{tX'}g)Q = \psi'_t(gQ),$$

and $\psi_t^0(gQ) = \psi_t(gQ)$. Hence $\{\psi^s\}$ is a homotopy consisting of $U(1)$ -actions between ψ and ψ' . $\square$

By $G_k(\mathbb{C}^n)$ we denote the Grassmann manifold consisting of the k -subspaces in $\mathbb{C}^n$. If M is a coadjoint orbit of $SU(n)$ diffeomorphic to $G_k(\mathbb{C}^n)$, it is the orbit of an element $\eta \in \mathfrak{su}(n)^*$ such that for $C \in \mathfrak{su}(n)$

$$(4.5) \quad \eta(C) = \alpha_1 \sum_{j=1}^k C_{jj} + \alpha_2 \sum_{j=k+1}^n C_{jj},$$

with $\alpha_1, \alpha_2 \in i\mathbb{R}$ and $\alpha_1 \neq \alpha_2$. That is, $\eta(C) = -ir \sum_{j=1}^k C_{jj}$, with $r \in \mathbb{R} \setminus \{0\}$.

Now we suppose that $k = 1$, then the coadjoint orbit $\mathcal{O} := \mathcal{O}_\eta = SU(n)/U(n-1) = \mathbb{C}P^{n-1}$. The coset $gU(n-1) \in SU(n)/U(n-1)$ determines the line in $\mathbb{C}^n$ defined by the first column of g . This is the identification $\mathcal{O} \simeq \mathbb{C}P^{n-1}$ we will use.

We write $\mathcal{R}$ for the set of roots determined by H , the diagonal subgroup of $SU(n)$. The stabilizer of η in the Weyl group $W = N(H)/H$ consists of all the permutations σ of $\{1, \dots, n\}$ with $\sigma(1) = 1$. Denoting by σ_1 the permutation identity and by σ_j the transposition $(1, j)$, with $j = 2, \dots, n$, then $\{\sigma_1, \dots, \sigma_n\}$ is a set of representatives of the classes in W/W_η .

Given $X \in \mathfrak{su}(n)$, the Hamiltonian isotopy generated by X has the map $h - \kappa$ as normalized Hamiltonian function, κ being a constant and h

$$(4.6) \quad h : \nu \in \mathcal{O} \mapsto -\langle \nu, X \rangle \in \mathbb{R}.$$

If $X = i(a_1, \dots, a_n)$ with $(a_j) \in (2\pi\mathbb{Z})^n$, then X generates a circle action ψ on $\mathcal{O}$. Furthermore, if $a_i \neq a_j$ for $i \neq j$, then the fixed point set for the action ψ is $\{\sigma_k \cdot \eta\}_{k=1, \dots, n}$.

Denoting by α_{ij} the root defined by $\alpha_{ij}(B) = b_i - b_j$, for $B = \text{diag}(b_k)$, then the set of positive roots is $\mathcal{R}^+ = \{\alpha_{12}, \dots, \alpha_{1n}\}$, assumed $r > 0$. For $k \neq 1$

$$(-i\sigma_k \cdot \alpha_{1s})(X) = \begin{cases} a_k - a_s, & \text{if } s \neq k \\ a_k - a_1, & \text{if } s = k \end{cases}.$$

And $(-i\sigma_1 \cdot \alpha_{1s})(X) = a_1 - a_s$. By (4.2) we obtain for the equivariant Euler class of normal bundle N_k to $\sigma_k \cdot \eta$

$$(4.7) \quad E(N_k) = \prod_{s \neq k} (a_k - a_s) u^{n-1}.$$

Since $h(\sigma_k \cdot \eta) = -\eta(\sigma_k^{-1} \cdot X) = -ra_k$, it follows from (4.7) together with (2.1)

$$(4.8) \quad S_\psi = \frac{e^{\kappa u}}{u^{n-1}} \sum_{k=1}^n \frac{e^{ra_k u}}{\prod_{s \neq k} (a_k - a_s)}.$$

Proposition 36. *Let $X = (x_1, \dots, x_n)$, $X' = (x'_1, \dots, x'_n)$ be regular elements of the integer lattice of $\mathfrak{h}$, and ψ, ψ' respective the circle actions on the coadjoint orbit $\mathcal{O}$ of $SU(n)$ diffeomorphic to $\mathbb{C}P^{n-1}$. Then the following statements are equivalent:*

- (a) $S_\psi = S_{\psi'}$.
- (b) *There is a permutation τ of $\{1, \dots, n\}$ such that $x'_j = x_{\tau(j)}$, for all j .*
- (c) $\psi \sim \psi'$.

Proof. (a) $\implies$ (b). If we write $X' = i(a'_1, \dots, a'_n)$ the expression for $S_{\psi'}$ can be obtained from (4.8) by substituting the a_j 's by the a'_j 's and κ by κ' . From the equality $S_\psi = S_{\psi'}$ together with the $\mathbb{R}[u, u^{-1}]$ -linear independence of the exponentials, it follows that there is a permutation τ of $\{1, \dots, n\}$ such that $\kappa' + ra'_j = \kappa + ra_{\tau(j)}$. Thus there is a constant c such that $a'_j - a_{\tau(j)} = c$ for all $j = 1, \dots, n$. As $\sum_j a'_j = \sum_i a_i = 0$, it turns out $c = 0$; hence (b) holds.

(b) $\implies$ (c) by Corollary 10. The equivalence of (a), (b) and (c) follows from Theorem 1. $\square$

Theorem 11 is the generalization of Proposition 36 when the hypothesis about the regularity of X and X' is deleted.

Proof of Theorem 11.

If

$$(4.9) \quad -iX = (\overbrace{a, \dots, a}^m, \overbrace{b, \dots, b}^m, \dots, \overbrace{d, \dots, d}^m) \equiv (a_1, \dots, a_n) \in (2\pi\mathbb{Z})^n$$

with $a, b, \dots$ different numbers, then a connected component of the fixed point set for the circle action ψ determined by X is $F = \{[z_1 : \dots : z_m : 0 : \dots : 0]\} \subset \mathbb{C}P^{n-1} = \mathcal{O}$. The contribution of the component F to S_ψ can be calculated as in the paragraphs before Lemma 19. That is, N_F can be decomposed in a direct sum of $U(1)$ -equivariant line bundles $\oplus_{j=m+1}^n N_j$, and

$$E(N_F) = \prod_{j=m+1}^n (c_1(N_j) + (a - a_j)u).$$

So we have as in (2.8)

$$(E(N_F))^{-1} = \frac{1}{u^{n-m}} \left(\alpha_0 + \frac{\alpha_1}{u} + \dots \right).$$

The aforementioned contribution to S_ψ is

$$e^{(\kappa+ra)u}Q_a(u),$$

$Q_a(u)$ being a Laurent polynomial of degree $m - n$. Thus expression for S_ψ is

$$(4.10) \quad S_\psi = e^{(\kappa+ra)u}Q_a(u) + \dots + e^{(\kappa+rd)u}Q_d(u).$$

Let

$$(4.11) \quad -iX' = (\overbrace{a', \dots, a'}^{m'}, \overbrace{e', \dots, e'}) \equiv (a'_1, \dots, a'_n) \in (2\pi\mathbb{Z})^n.$$

The equality $S_\psi = S_{\psi'}$ implies the following set equality $\{a + \beta, b + \beta, \dots, d + \beta\} = \{a', \dots, e'\}$, with β constant. Furthermore, if for example $a' = b + \beta$ then $\text{degree } Q_{a'} = \text{degree } Q_b$. That is, the number of b 's in (4.9) must be equal to the number of a' 's in (4.11). Hence there is a permutation τ of $\{1, \dots, n\}$, such that $a_j + \beta = a'_{\tau(j)}$ and if $a_i = a_j$ then $a'_{\tau(i)} = a'_{\tau(j)}$. As $\sum_j a_j = 0 = \sum_i a'_i$, then $\beta = 0$. So the theorem is consequence of Corollary 10 and Theorem 1. $\square$

Proof of Theorem 12.

Now we must consider the case $1 < k < n$ in (4.5). In this case, if the permutation $\sigma \in W_\eta$, then σ applies the subset $\{1, \dots, k\}$ onto itself (and of course $\{k+1, \dots, n\}$ onto itself). The stabilizer W_η has $k!(n-k)!$ elements. A set of representatives of the elements of W/W_η is the set $\mathcal{C}_k$ of combinations of $\{1, \dots, n\}$ with $n-k$ elements. If $\mathfrak{d} \in \mathcal{C}_k$, $\sigma_{\mathfrak{d}}$ will denote the corresponding element in the quotient W/W_η .

Let $X = \text{diag}(ia_1, \dots, ia_n) \in (2\pi i\mathbb{Z})^n$ be a regular element, and ψ is the corresponding circle action on the Grassmannian $G_k(\mathbb{C}^n)$. If $\mathfrak{c} = \{1, \dots, n\} \setminus \{j_1, \dots, j_k\}$, the value of the Hamiltonian (4.6) on $\mathfrak{c} \cdot \eta$ is

$$h(\sigma_{\mathfrak{c}} \cdot \eta) = -r \sum_{l=1}^k a_{j_l}.$$

We write $\alpha_{\mathfrak{d}} = \sum_{l=1}^k a_{i_l}$, when $\mathfrak{d} = \{i_1, \dots, i_k\} \in \mathcal{C}_k$. Then

$$(4.12) \quad S_\psi = \frac{e^{\kappa u}}{u^{k(n-k)}} \sum_{\mathfrak{c} \in \mathcal{C}_k} \frac{e^{r\alpha_{\mathfrak{c}} u}}{p_{\mathfrak{c}}},$$

where $p_{\mathfrak{c}} \in \mathbb{Z}$ and r is a real number.

If X' is another regular element and $S_\psi = S_{\psi'}$, it follows from (4.12) that there is a bijective map $f : \mathcal{C}_k \rightarrow \mathcal{C}_k$ such that

$$\kappa + \alpha_{f(\mathfrak{c})} = \kappa' + \alpha'_{\mathfrak{c}},$$

for all $\mathfrak{c} \in \mathcal{C}_k$. $\square$

If $\mathcal{O}$ is a coadjoint orbit of $SU(n)$ diffeomorphic to the full flag manifold, it is the orbit of an element $\eta \in \mathfrak{su}(n)^*$, such that for any $C \in \mathfrak{su}(n)$

$$(4.13) \quad \eta(C) = -i \sum_{j=1}^{n-1} r_j C_{jj},$$

with $r_i \neq r_j$ for $i \neq j$. We will assume that $r_1 > r_2 > \dots > r_{n-1}$. So $\mathcal{R}^+ = \{\alpha_{ij} : i < j\}$.

Given $X \in \mathfrak{su}(n)$ with $-iX = (a_1, \dots, a_n) \in (2\pi\mathbb{Z})^n$ and $a_i \neq a_j$ for $i \neq j$, the fixed point set for the S^1 -action ψ defined by X on $\mathcal{O}$ is $\{\sigma \cdot \eta : \sigma \in \mathcal{S}_n\}$, where $\mathcal{S}_n$ is the symmetric group. The $U(1)$ -equivariant Euler class

$$E(N_{\sigma \cdot \eta}) = u^m \epsilon(\sigma) \prod_{r < s} (a_r - a_s),$$

$\epsilon(\sigma)$ being the signature of σ , and $m = n(n-1)/2$. As

$$-\eta(\sigma \cdot X) = \sum_{j=1}^{n-1} r_j a_{\sigma(j)},$$

we obtain

$$(4.14) \quad S_\psi = \frac{e^{\kappa u}}{u^m \prod_{r < s} (a_r - a_s)} \sum_{\sigma \in \mathcal{S}_n} \epsilon(\sigma) \exp\left(u \sum_{j=1}^{n-1} r_j a_{\sigma(j)}\right).$$

If X' is another regular element with $-iX' = (a'_1, \dots, a'_n) \in (2\pi\mathbb{Z})^n$ and $S_\psi = S_{\psi'}$, it follows from (4.14) together with Lemma 14 that there exist a constant β and a bijective map f from $\mathcal{S}_n$ into itself, such that for all $\sigma \in \mathcal{S}_n$

$$(4.15) \quad \sum_{j=1}^{n-1} r_j a'_{\sigma(j)} = \sum_{j=1}^{n-1} r_j a_{f\sigma(j)} + \beta.$$

We have proved the following Proposition

Proposition 37. *Let $\mathcal{O}$ be the coadjoint orbit of the element $\eta \in \mathfrak{su}(n)^*$ given by (4.13), and let ψ and ψ' be the $U(1)$ -actions defined by the regular elements $(ia_1, \dots, ia_n)$ and $(ia'_1, \dots, ia'_n)$ respectively. If $\psi \sim \psi'$, then there exist a constant β and a bijective map f from $\mathcal{S}_n$ into itself, such that (4.15) holds for all $\sigma \in \mathcal{S}_n$.*

Remark. In particular, if there exist $\tau \in \mathcal{S}_n$, such that $a'_j = a_{\tau(j)}$ for all j ; that is, the hypothesis of Corollary 10 are satisfied, then we can construct the map $f : \sigma \in \mathcal{S}_n \mapsto \tau \cdot \sigma \in \mathcal{S}_n$. Thus $a'_{\sigma(k)} = a_{f\sigma(k)}$ and equation (4.15) holds with $\beta = 0$. That is, Proposition 37 is consistent with Corollary 10.

REFERENCES

- [1] Abreu, M., McDuff, D. Topology of symplectomorphism groups of rational ruled surfaces. J. Amer. Math. Soc. **13** 971-1009 (2000).
- [2] Atiyah, M. Convexity and commuting Hamiltonians. Bul. London Math. Soc. **14**, 1-15 (1982)
- [3] Atiyah, M. F., Bott, R.: The moment map and equivariant cohomology. Topology **23**, 1-28 (1984)
- [4] Berline, N., Getzler, E., Vergne, M.: Heat Kernels and Dirac Operators. Springer-Verlag, Berlin (1991)
- [5] Delzant, T.: Hamiltoniens périodiques et image convexe de l'application moment. Bull. Soc. Math. France **116**, 315-339 (1988)
- [6] Fulton, W., Harris, J.: Representation Theory. Springer-Verlag, New York (1991)
- [7] Goodman, R., Wallach, N. R.: Representations and Invariants of Classical Groups. Cambridge U. P., Cambridge (1998)
- [8] Gromov, M.: Pseudo holomorphic curves in symplectic manifolds. Invent. Math. **82**, 307-347 (1985)
- [9] Guillemin, V.: Moment maps and combinatorial invariants of Hamiltonian T^n -spaces. Birkhäuser, Boston, (1994)
- [10] Guillemin, V., Lerman, L., Sternberg, S.: Symplectic fibrations and multiplicity diagrams. Cambridge U.P., Cambridge (1996)

- [11] Guillemin, V., Sternberg, S.: Convexity properties of the moment mapping. *Invent. Math.* **67**, 491-513 (1982)
- [12] Guillemin, V., Sternberg, S.: *Symplectic techniques in Physics*. Cambridge U.P., Cambridge (1984)
- [13] Guillemin, V., Sternberg, S.: *Supersymmetry and equivariant de Rham theory*. Springer-Verlag, Berlin (1999)
- [14] Januszkiewicz, T., Kedra, J.: Characteristic classes of smooth fibrations. <http://arXiv.org/math/0209288>
- [15] Karshon, Y.: Maximal Tori in the Symplectomorphism Groups of Hirzebruch Surfaces. *Math. Res. Lett.* **10**, 125-132 (2003)
- [16] Kedra, J., McDuff, D.: Homotopy properties of Hamiltonian group actions. *Geom. Topol.* **9**, 121-162 (2005)
- [17] Kirilov, A.A.: *Elements of the Theory of Representations*. Springer-Verlag, Berlin (1976)
- [18] Lerman, E., Tolman S.: Hamiltonian torus actions on symplectic orbifolds and toric varieties. *Trans. Amer. Math. Soc.* **10**, (1997) 4201-4230 (1997)
- [19] McDuff, D. Symplectomorphism groups and almost complex structures. *Monogr. Enseign. Math.* **38** (2001) ([arXiv:math.SG/0010274](http://arXiv.org/math.SG/0010274))
- [20] McDuff, D., Salamon, D.: *Introduction to symplectic topology*. Clarenton Press, Oxford (1998)
- [21] Polterovich, L.: *The geometry of the group of symplectic diffeomorphisms*. Birkhäuser, Basel, (2001)
- [22] Viña, A.: A characteristic number of Hamiltonian bundles over S^2 . *J. Geom. Phys.* **56**, 2327-2343 (2006)
- [23] Viña, A.: Hamiltonian diffeomorphisms of toric manifolds and flag manifolds. *J. Geom. Phys.* **57**, 943-965 (2007)

DEPARTAMENTO DE FÍSICA. UNIVERSIDAD DE OVIEDO. AVDA CALVO SOTELO. 33007 OVIEDO. SPAIN.

E-mail address: vina@uniovi.es